

THE HOMOCLINIC AND HETEROCLINIC C^* -ALGEBRAS OF A GENERALIZED ONE-DIMENSIONAL SOLENOID

KLAUS THOMSEN

1. INTRODUCTION

The homoclinic and heteroclinic structure in dynamical systems was first used to produce C^* -algebras in the way breaking work of Cuntz and Krieger in [Kr1], [Kr2] and [CuK]. This work has been generalized in many directions where the relation to dynamical systems is either absent or appears very implicit, but Ian Putnam described in [Pu] a natural way to extend the constructions of Cuntz and Krieger to higher dimensions such that the point of departure is the heteroclinic structure in a Smale space, just as the work of Cuntz and Krieger departed from the heteroclinic structure in a shift of finite type, which is a zero-dimensional Smale space. Putnam builds his approach on the work of D. Ruelle, [Ru1], [Ru2], who introduced the notion of a Smale space in [Ru1] and constructed the so-called asymptotic algebra from the homoclinic equivalence relation in [Ru2].

The work of Putnam and Ruelle was further generalized by the author in [Th1] where it was shown that Ruelle's approach can be adopted as soon as there is enough expansiveness in the underlying dynamical system; the local product structure in a Smale space is not crucial for the construction. Furthermore, in [Th1] the alternative approach was used to obtain inductive limit decompositions for the algebras of Putnam arising from particular classes of Smale spaces, e.g. expansive group automorphisms and one-dimensional generalized solenoids in the sense of R.F. Williams, [Wi2], and I. Yi, [Y1]. For expansive group automorphisms it was shown that the C^* -algebras are all AT-algebras of real rank zero, and hence are classified by their K-theory groups, thanks to the work of G. Elliott, [Ell1]. For one-dimensional generalized solenoids the exact nature of the inductive limit decomposition was not determined and the homoclinic algebra was not examined. In particular, it was not decided if the C^* -algebras are classified by K-theory. The main purpose of the present paper is tie up this loose end by showing that they are, although they turn out to be more general AH-algebras and exhibit more complicated K-theory than the algebras arising from expansive group automorphisms, at least in the sense that torsion appears. Specifically, it is shown that the heteroclinic algebra of both a one-solenoid and its inverse, as well as the homoclinic algebra are all AH-algebras of real rank zero with no dimension growth. They are therefore classified by K-theory thanks to the work of Elliott and Gong, [EG]. This conclusion is obtained for the heteroclinic algebra by combining a thorough study of the inductive limit decomposition obtained in [Th1] with results on the classification of simple C^* -algebras, in particular results by H. Lin on algebras of tracial rank zero, cf. e.g. [Lin4].

The study of the K_1 -group of the heteroclinic algebra reveals a fundamental dichotomy between the orientable and the non-orientable case. For orientable one-solenoids the K_1 -group of the heteroclinic algebra is $\mathbb{Z}$ while it collapses to $\mathbb{Z}_2$ when the one-solenoid is not orientable. It follows from this that a one-solenoid is only conjugate to an orientable one-solenoid when the graph map defining it is already oriented.

The dichotomy between the oriented and non-oriented case turns out to be of pervasive importance for the structure of the three algebras arising from the heteroclinic and homoclinic structure of a one-solenoid. For example, one of these algebras is an AT-algebra if and only if they all are, and this happens if and only if the one-solenoid is orientable. In the oriented case the heteroclinic algebra of a one-solenoid is not so different from the heteroclinic algebra of its inverse; they are both AT-algebras and have the same K_1 -group, namely $\mathbb{Z}$. In fact the two algebras are often, but not always, isomorphic in this case. In contrast the heteroclinic algebra of the inverse of a one-solenoid differs substantially from the heteroclinic algebra of the one-solenoid itself in the non-orientable case. It turns out that the heteroclinic algebra of the inverse of a one-solenoid is the stabilization of a crossed product C^* -algebra coming from a free and minimal action of the infinite dihedral group on the Cantor set, and the results on its structure are obtained through a study of such crossed products. Again the work of Lin on algebras with tracial rank zero plays a fundamental role, one of the crucial steps being that we can use results of N.C. Phillips on finite group actions with the tracial Rokhlin property, [Ph4],[Ph5], to conclude that the crossed product C^* -algebra coming from a free and minimal action of the infinite dihedral group on the Cantor set has tracial rank zero. Among the results we obtain is that the K_1 -group vanishes for the heteroclinic algebra of the inverse of a non-orientable one-solenoid, but nonetheless the algebra is not an AF-algebra because the two-torsion now pops up in the K_0 -group.

The results we obtain on the structure of the homoclinic algebra of a one-solenoid is obtained from the results on the heteroclinic algebras by use of a result of Ian Putnam from [Pu] which says that the homoclinic algebra is stably isomorphic to the crossed product of the two heteroclinic algebras.

Acknowledgement: I want to thank Marcy Barge and Ian Putnam for their help with this paper. In particular, I thank Marcy for convincing me that an unorientable one-solenoid is a Smale space, and Ian for many discussions and for pointing out a mistake in the first version which concealed the role of the infinite dihedral group.

2. THE DYNAMICAL SETUP

2.1. The homoclinic and heteroclinic algebras of an expansive homeomorphism. Let (X, d) be a compact metric space and $\psi : X \rightarrow X$ a homeomorphism. Recall that ψ is *expansive* when there is a $\delta > 0$ such that

$$x \neq y \Rightarrow \sup_{k \in \mathbb{Z}} d(\psi^k(x), \psi^k(y)) \geq \delta.$$

We say that two points $x, y \in X$ are *locally conjugate* when there are open neighborhoods U and V of x and y , respectively, and a homeomorphism $\chi : U \rightarrow V$ such that $\chi(x) = y$ and $\lim_{k \rightarrow \pm\infty} \sup_{z \in U} d(\psi^k(z), \psi^k(\chi(z))) = 0$. This is an equivalence relation on X and

its graph

$$R_\psi(X) = \{(x, y) \in X^2 : x \text{ is locally conjugate to } y\}$$

is a locally compact Hausdorff space in the topology for which every local conjugacy (U, V, χ) defines an element of a canonical base:

$$\{(z, \chi(z)) : z \in U\}.$$

This topology is typically different from the topology which $R_\psi(X)$ inherits from $X \times X$; it has more open sets. The crucial fact is that $R_\psi(X)$ is what is nowadays known as an étale equivalence relation, cf. [Ph1], [GPS1], [Th1], so that the reduced groupoid C^* -algebra $C_r^*(R_\psi(X))$ of Renault, [Re], can be defined. We call this *the homoclinic algebra* of (X, ψ) and denote it by $A_\psi(X)$. When (X, ψ) is a Smale space, as defined by Ruelle in [Ru1], the homoclinic algebra $A_\psi(X)$ is the asymptotic algebra of Ruelle and Putnam, [Ru2],[Pu]. When X is a sub-shift the homoclinic algebra is the AF-algebra whose dimension group was defined by Krieger in [Kr1] and Section 2 of [Kr2].

Let $\text{Per } \psi$ denote the set of ψ -periodic points. We assume that $\text{Per } \psi \neq \emptyset$. For $p \in \text{Per } \psi$, set

$$W^u(p) = \{x \in X : \lim_{k \rightarrow -\infty} d(\psi^k(x), \psi^k(p)) = 0\}.$$

Since ψ is expansive each $W^u(p)$ is a locally compact Hausdorff space in a topology with base

$$\{y \in X : d(\psi^j(y), \psi^j(x)) < \epsilon, j \leq k\},$$

where $x \in W^u(p)$, $k \in \mathbb{Z}$, $\epsilon \in]0, \epsilon_p[$ are arbitrary, and $\epsilon_p > 0$ only depends on p . See Lemma 4.6 of [Th1]. The set of *post-periodic points* of (X, ψ) is

$$W_{X, \psi} = \bigcup_{p \in \text{Per } \psi} W^u(p),$$

and it is a locally compact Hausdorff space in the topology, which we call *the Wagoner topology*, defined such that each $W^u(p)$ is open in $W_{X, \psi}$ and has the topology we have just described above.

Define the equivalence relation $\sim$ on $W_{X, \psi}$ such that $x \sim y$ if and only if there are open neighborhoods, U of x and V of y in $W_{X, \psi}$, and a homeomorphism $\gamma : U \rightarrow V$, again called *a local conjugacy*, such that $\gamma(x) = y$ and

$$\lim_{k \rightarrow \infty} \sup_{z \in U} d(\psi^k(z), \psi^k(\gamma(z))) = 0.$$

This is an étale equivalence relation $R_\psi(X, W_{X, \psi})$, cf. [Th1], and the corresponding (reduced) groupoid C^* -algebra $C_r^*(R_\psi(X, W_{X, \psi}))$ is *the heteroclinic algebra* of (X, ψ) . As in [Th1] we denote it by $B_\psi(X)$. When (X, ψ) is a mixing Smale space the heteroclinic algebra is $*$ -isomorphic to the stabilization of Putnam's 'stable algebra', cf. [Pu] and Theorem 4.17 of [Th1]. In particular, it is a higher dimensional analogue of the 'AF-core' in the Cuntz-Krieger construction, cf. [CuK].

2.2. Generalized one-dimensional solenoids. Let Γ be a finite (unoriented) graph with vertexes $\mathbb{V}$ and edges $\mathbb{E}$. Consider a continuous map $h : \Gamma \rightarrow \Gamma$ such that the following conditions are satisfied for some metric d for the topology of Γ :

α) (Expansion) There are constants $C > 0$ and $\lambda > 1$ such that

$$d(h^n(x), h^n(y)) \geq C\lambda^n d(x, y)$$

for every $n \in \mathbb{N}$ when $x, y \in e \in \mathbb{E}$ and there is an edge $e' \in \mathbb{E}$ with $h^n([x, y]) \subseteq e'$.
 ($[x, y]$ is the interval in e between x and y .)

β) (Non folding) h^n is locally injective on e for each $e \in \mathbb{E}$ and each $n \in \mathbb{N}$.

γ) (Markov) $h(\mathbb{V}) \subseteq \mathbb{V}$.

δ) (Mixing) For every edge $e \in \mathbb{E}$ there is an $m \in \mathbb{N}$ such that $\Gamma \subseteq h^m(e)$.

ϵ) (Flattening) There is a $d \in \mathbb{N}$ such that for all $x \in \Gamma$ there is a neighborhood U_x of x with $h^d(U_x)$ homeomorphic to $] -1, 1[$.

When all conditions hold we are in a setting first introduced by Williams in [Wi2] and later studied by I. Yi in [Y1],[Y2],[Y3],[Y4],[Y5]. We say then that (Γ, h) is a *pre-solenoid*.¹

Set

$$\bar{\Gamma} = \{(x_i)_{i=0}^\infty \in \Gamma^\mathbb{N} : h(x_{i+1}) = x_i, i = 0, 1, 2, \dots\}.$$

We consider $\bar{\Gamma}$ as a compact metric space with the metric

$$D((x_i)_{i=0}^\infty, (y_i)_{i=0}^\infty) = \sum_{i=0}^\infty 2^{-i} d(x_i, y_i).$$

Define $\bar{h} : \bar{\Gamma} \rightarrow \bar{\Gamma}$ such that $\bar{h}(x)_i = h(x_i)$ for all $i \in \mathbb{N}$. $\bar{h}$ is a homeomorphism with inverse

$$\bar{h}^{-1}(z_0, z_1, z_2, \dots) = (z_1, z_2, z_3, \dots).$$

Following Williams and Yi, [Wi2], [Y1], we call $(\bar{\Gamma}, \bar{h})$ a *generalized one-dimensional solenoid* or just a *1-solenoid*.

It was shown by Williams in [Wi1] that expanding attractors of certain diffeomorphisms of compact manifolds are 1-solenoids via a conjugacy which turns the restriction of the diffeomorphism into $\bar{h}$, and that each 1-solenoid arises in this way from a diffeomorphism of the 4-sphere.

As we shall see there is a dichotomy in the class of 1-solenoids which is crucial for the structure of the C^* -algebras they give rise to and which depends on whether or not the 1-solenoid is orientable. To formalize this notion we define an *orientation* of Γ to be a collection of homeomorphisms $\psi_e : [0, 1] \rightarrow e$, $e \in \mathbb{E}$. We say that h is *positively oriented* (resp. *negatively oriented*) with respect to the orientation $\psi_e, e \in \mathbb{E}$, when the function

$$\psi_{e'}^{-1} \circ h \circ \psi_e : \psi_e^{-1}(e \cap h^{-1}(e')) \rightarrow [0, 1] \quad (2.1)$$

is increasing (resp. decreasing) for every pair $e, e' \in \mathbb{E}$. A pre-solenoid (Γ, h) is *positively oriented* (resp. *negatively oriented*) when there is an orientation of the edges in Γ such that h is positively oriented (resp. negatively) oriented with respect to that orientation. (Γ, h) is *oriented* when it is either positively or negatively oriented.

¹Unlike what I believed when [Th1] was written, the combined conditions are strictly stronger than the conditions considered by Williams and Yi. If for example Γ consists of two disjoint circles and h takes one circle twice around the other in an appropriate expanding way (like $z \mapsto z^2$ on the unit circle in the complex plane), then δ) fails, but all the Axioms 0-5 of [Y1] are satisfied. The example shows that Lemma 2.14 of [Y1] is wrong. The mistake occurs in the proof of 1.6 Lemma in [Wi2].

The 1-solenoid $(\bar{\Gamma}, \bar{h})$ is *orientable* where there is an oriented pre-solenoid (Γ_1, h_1) such that $(\bar{\Gamma}, \bar{h})$ is conjugate to $(\bar{\Gamma}_1, \bar{h}_1)$. When (Γ_1, h_1) can be chosen to be positively (resp. negatively) oriented we say that $(\bar{\Gamma}, \bar{h})$ is *positively (resp. negatively) orientable*.

Remark 2.1. Our terminology concerning orientation of pre-solenoids is more elaborate than that used by Yi, cf. [Y1]-[Y5]. To illustrate the difference, let Γ be the wedge of two circles a and b , and let $h : \Gamma \rightarrow \Gamma$ be given by the wrapping rule

$$a \mapsto a^{-1}b^{-1}, \quad b \mapsto a^{-1}b^{-1}$$

in the sense of Definition 2.6 in [Y1]. Then (Γ, h) is a pre-solenoid which is negatively oriented but not positively oriented. In contrast, the wrapping rule

$$a \mapsto ba, \quad b \mapsto ba$$

is positively oriented and not negatively oriented. Despite the fact that the square of the two maps are identical and positively oriented, the 1-solenoids they define are not conjugate.

Through the study of the heteroclinic C^* -algebra $B_{\bar{h}}(\bar{\Gamma})$ of a 1-solenoid $(\bar{\Gamma}, \bar{h})$, in particular a careful examination of its K_1 -group, we shall obtain the following result.

Theorem 2.2. *Let (Γ, h) be a pre-solenoid. Then $(\bar{\Gamma}, \bar{h})$ is positively orientable (resp. negatively orientable) if and only if (Γ, h) is positively oriented (resp. negatively oriented).*

3. THE HETEROCLINIC ALGEBRA OF A 1-SOLENOID

In this section we study the structure of the heteroclinic algebra of a 1-solenoid. The point of departure is the inductive limit decomposition of the heteroclinic algebra obtained in Theorem 5.17 of [Th1].

3.1. The building blocks. Let Γ be a finite graph and $h : [-1, 1] \rightarrow \Gamma$ a locally injective continuous map. We define an equivalence relation $\sim$ on $] - 1, 1[$ such that $t \sim s$ if and only if $h(t) = h(s)$ and there are open neighborhoods U_s and U_t of s and t in $] - 1, 1[$, respectively, such that $h(U_s) = h(U_t) \simeq] - 1, 1[$. Set

$$R_h = \{(s, t) \in] - 1, 1[^2 : s \sim t\}.$$

Give R_h the topology inherited from $] - 1, 1[^2$.

Lemma 3.1. *R_h is an étale equivalence relation.*

Proof. This is part of Lemma 5.13 in [Th1]. □

We are going to use the étale equivalence relations of Lemma 3.1 in the special case where $h(-1), h(1) \in \mathbb{V}$ and $h(] - 1, 1]) = \Gamma$. When this holds we say that R_h is an *open interval-graph relation*.

To give a manageable description of $C_r^*(R_h)$, let $m, n \in \mathbb{N}$. Consider some $a = (a(1), a(2), \dots, a(m)) \in \mathbb{N}^m$, $b = (b(1), b(2), \dots, b(n)) \in \mathbb{N}^n$ and two $n \times m$ -matrices, I, U , with $\{0, 1\}$ -entries. Assume that

$$\sum_{k=1}^m I_{ik} a(k) \leq b(i) \tag{3.1}$$

and

$$\sum_{k=1}^m U_{ik} a(k) \leq b(i) \quad (3.2)$$

for all $i = 1, 2, \dots, n$. Set $F_a = M_{a(1)} \oplus M_{a(2)} \oplus \dots \oplus M_{a(m)}$ and $F_b = M_{b(1)} \oplus M_{b(2)} \oplus \dots \oplus M_{b(n)}$, where M_k denotes the C^* -algebra of complex $k \times k$ matrices. Let $\varphi^I, \varphi^U : F_a \rightarrow F_b$ be $*$ -homomorphisms with partial maps $\varphi_{ij}^I, \varphi_{ij}^U : M_{a(j)} \rightarrow M_{b(i)}$ of multiplicity I_{ij} and U_{ij} , respectively. Set

$$A(a, b, I, U) = \{(x, f) \in F_a \oplus C([0, 1], F_b) : \varphi^I(x) = f(0), \varphi^U(x) = f(1)\}.$$

In the following we shall only be interested in algebras of this type when

$$\sum_{i=1}^n I_{ik} + \sum_{i=1}^n U_{ik} = 2 \quad (3.3)$$

for all $k = 1, 2, \dots, m$. We will refer to such an algebra as a *building block*. Note that $A(a, b, I, U)$ is unital if and only if

$$\sum_{k=1}^m U_{ik} a(k) = \sum_{k=1}^m I_{ik} a(k) = b(i)$$

for all i .

For the explicit identification of $C_r^*(R_h)$ with a building block we need to decide for each edge which of its endpoints corresponds to $0 \in [0, 1]$ and which to $1 \in [0, 1]$. For this purpose we give each edge of Γ an (arbitrary) orientation so that they may be considered as directed arrows instead of undirected edges. The start vertex of the arrow e is then denoted by $s(e)$ and the terminal vertex of e by $t(e)$. For each edge e of Γ choose a homeomorphism $\psi_e : e \rightarrow [0, 1]$ such that $\psi_e(s(e)) = 0$, $\psi_e(t(e)) = 1$.

Let $-1 = x_0 < x_1 < x_2 < \dots < x_N = 1$ be the elements of $h^{-1}(\mathbb{V})$. A *passage* in Γ is a closed subset $J \subseteq \Gamma$ containing exactly one vertex v such that there is a homeomorphism $\varphi : J \rightarrow [-1, 1]$ with $\varphi^{-1}(0) = v$. We will identify two passages J and J' in Γ when $J \cap J'$ is also a passage in Γ . A *h-passage* is a passage in Γ which contains $h([x_i - \epsilon, x_i + \epsilon])$ for some $i \in \{1, 2, \dots, N - 1\}$ and all sufficiently small $\epsilon > 0$. An arrow e is then an *entry-arrow* in the h -passage p when $s(e) \in p$ and an *exit-arrow* when $t(e) \in p$.

Note that for every element $x \in h^{-1}(\mathbb{V})$ there is a unique h -passage p_x such that $p_x = h([x - \epsilon, x + \epsilon])$ for all small $\epsilon > 0$. Let $\mathbb{A}_h$ denote the finite-dimensional C^* -algebra generated by the matrix-units $e_{x,y}$, $x, y \in h^{-1}(\mathbb{V})$, such that $p_x = p_y$. Similarly, let $\mathcal{I}_h$ denote the set of connected components of $]0, 1[\setminus h^{-1}(\mathbb{V})$ and let $\mathbb{B}_h$ denote the finite-dimensional C^* -algebra generated by the matrix units $e_{I,J}$, where $I, J \in \mathcal{I}_h$ and $h(\overline{I}) = h(\overline{J})$. Define $\pi^I : \mathbb{A}_h \rightarrow \mathbb{B}_h$ such that $\pi^I(e_{x,y}) = \sum_{J,K} e_{J,K}$ where we sum over the pairs $J, K \in \mathcal{I}_h$ with the property that $x \in \overline{J}, y \in \overline{K}, h(\overline{J}) = h(\overline{K})$ and $h(\overline{J})$ is an entry-arrow in both p_x and p_y . There are at most two such pairs - when there is none we set $\pi^I(e_{x,y}) = 0$. Define $\pi^U : \mathbb{A}_h \rightarrow \mathbb{B}_h$ such that $\pi^U(e_{x,y}) = \sum_{J,K} e_{J,K}$ where we sum over the pairs $J, K \in \mathcal{I}_h$ with the property that $x \in \overline{J}, y \in \overline{K}, h(\overline{J}) = h(\overline{K})$ and $h(\overline{J})$ is an exit-arrow in both p_x and p_y . There are at most two such pairs - when there is none we set $\pi^U(e_{x,y}) = 0$. Set

$$\mathbb{D}_h = \{(x, f) \in \mathbb{A}_h \oplus C([0, 1], \mathbb{B}_h) : f(0) = \pi^I(x), f(1) = \pi^U(x)\}.$$

We can then define a $*$ -isomorphism $\Phi_h : C_r^*(R_h) \rightarrow \mathbb{D}_h$ such that

$$\Phi_h(f) = \left(\sum_{x,y \in h^{-1}(\mathbb{V})} f(x,y) e_{x,y}, \sum_{I,J \in \mathcal{I}_h} f\left((h|_I)^{-1} \circ \psi_h^{-1}(\cdot), (h|_J)^{-1} \circ \psi_h^{-1}(\cdot)\right) e_{I,J} \right)$$

when $f \in C_c(R_h)$. This is essentially the $*$ -isomorphism from Lemma 5.16 of [Th1], revised to avoid the assumption about the absence of loops in Γ which was imposed there.

Let $p_i, i = 1, 2, \dots, m$, be a numbering of the different h -passages and $e_i, i = 1, 2, \dots, n$, a numbering of the edges in Γ . Set

$$a(i) = \# \{j \in \{1, 2, \dots, N-1\} : h([x_j - \epsilon, x_j + \epsilon]) = p_i \text{ for all small } \epsilon\},$$

$i = 1, 2, \dots, m$, and

$$b(i) = \# \{j \in \{0, 1, \dots, N-1\} : h([x_j, x_{j+1}]) = e_i\},$$

$i = 1, 2, \dots, n$. Set $a = (a(1), a(2), \dots, a(m)) \in \mathbb{N}^m$, $b = (b(1), b(2), \dots, b(n)) \in \mathbb{N}^n$. Then there are obvious $*$ -isomorphisms $\kappa : \mathbb{A}_h \rightarrow F_a$ and $\kappa' : \mathbb{B}_h \rightarrow F_b$. Let

$$I_{ik} = \begin{cases} 1, & \text{when } e_i \text{ is an entry-edge in } p_k \\ 0, & \text{otherwise,} \end{cases}$$

and

$$U_{ik} = \begin{cases} 1, & \text{when } e_i \text{ is an exit-edge in } p_k \\ 0, & \text{otherwise.} \end{cases}$$

Note that (3.3) holds, i.e. $A(a, b, I, U)$ is a building block. Furthermore, $\varphi^I = \kappa' \circ \pi^I \circ \kappa^{-1}$ and $\varphi^U = \kappa' \circ \pi^U \circ \kappa^{-1}$, at least up to unitary equivalences which we can safely ignore. We get therefore a $*$ -isomorphism $\Psi_h : \mathbb{D}_h \rightarrow A(a, b, I, U)$ defined such that

$$\Psi_h(x, f) = (\kappa(x), (\text{id}_{C[0,1]} \otimes \kappa')(f)).$$

In this way we get

Lemma 3.2. (cf. Lemma 5.16 of [Th1]) $\Psi_h \circ \Phi_h : C_r^*(R_h) \rightarrow A(a, b, I, U)$ is a $*$ -isomorphism.

Note that the building block $A(a, b, I, U)$ of Lemma 3.2 satisfies condition (3.4) of the following lemma by construction.

Lemma 3.3. Let $A(a, b, I, U)$ be a building block corresponding to the vectors $a = (a(1), a(2), \dots, a(m)) \in \mathbb{N}^m$ and $b = (b(1), b(2), \dots, b(n)) \in \mathbb{N}^n$. Assume that

$$\sum_{j=1}^m U_{kj} \geq 1, \quad \sum_{j=1}^m I_{kj} \geq 1 \quad (3.4)$$

for all $k = 1, 2, \dots, n$, and $a(i) \geq 2n + 1$ for all $i = 1, 2, \dots, m$. Then $A(a, b, I, U)$ contains a non-zero projection.

Proof. We define a labeled graph $\mathcal{G}$ as follows: The vertexes of $\mathcal{G}$ consist of the tuples $(i, \rightarrow), i = 1, 2, \dots, n$, and the tuples $(i, \leftarrow), i = 1, 2, \dots, n$. The arrows $\rightsquigarrow$ in $\mathcal{G}$ are labeled by the elements of $\{1, 2, \dots, m\}$ and there is a labeled arrow $(i, \rightarrow) \xrightarrow{j} (i', \rightarrow)$ when $\varphi_{ij}^U \neq 0$ and $\varphi_{i'j}^I \neq 0$, there is a labeled arrow $(i, \rightarrow) \xrightarrow{j} (i', \leftarrow)$ when $i \neq i', \varphi_{ij}^U \neq 0$

and $\varphi_{i'j}^U \neq 0$, there is a labeled arrow $(i, \leftarrow) \xrightarrow{j} (i', \leftarrow)$ when $\varphi_{ij}^I \neq 0$ and $\varphi_{i'j}^U \neq 0$ and there is a labeled arrow $(i, \leftarrow) \xrightarrow{j} (i', \rightarrow)$ when $i \neq i'$, $\varphi_{ij}^I \neq 0$ and $\varphi_{i'j}^I \neq 0$.

Given an arrow $(i, \rightarrow) \xrightarrow{j} (i', \rightarrow)$ in $\mathcal{G}$ we write $p \xrightarrow{q} p'$ when $p \in C([0, 1], M_{b(i)})$, $p' \in C([0, 1], M_{b(i')})$ and $q \in M_{a(j)}$ are rank 1 projections such that $p(1) = \varphi_{ij}^U(q)$ and $\varphi_{i'j}^I(q) = p'(0)$. We say that $p \xrightarrow{q} p'$ realizes $(i, \rightarrow) \xrightarrow{j} (i', \rightarrow)$. In the following all projections, with a single obvious exception, will be rank 1 projections. We say that $p \xrightarrow{q} p'$ realizes $(i, \rightarrow) \xrightarrow{j} (i', \leftarrow)$ when $p(1) = \varphi_{ij}^U(q)$ and $\varphi_{i'j}^U(q) = p'(1)$, that $p \xrightarrow{q} p'$ realizes $(i, \leftarrow) \xrightarrow{j} (i', \leftarrow)$ when $p(0) = \varphi_{ij}^I(q)$ and $\varphi_{i'j}^U(q) = p'(1)$ and finally that $p \xrightarrow{q} p'$ realizes $(i, \leftarrow) \xrightarrow{j} (i', \rightarrow)$ when $p(0) = \varphi_{ij}^I(q)$ and $\varphi_{i'j}^I(q) = p'(0)$. Given a finite path

$$c = (i_1, *_1) \xrightarrow{j_1} (i_2, *_2) \xrightarrow{j_2} \dots \xrightarrow{j_{d-1}} (i_d, *_d)$$

in $\mathcal{G}$, where $*_k \in \{\leftarrow, \rightarrow\}$, we say that c is realized by $p_1 \xrightarrow{q_1} p_2 \xrightarrow{q_2} p_3 \xrightarrow{q_3} \dots \xrightarrow{q_{d-1}} p_d$ when $p_k \xrightarrow{q_k} p_{k+1}$ realizes $(i_k, *_{k+1}) \xrightarrow{j_{k+1}} (i_{k+1}, *_{k+1})$ for all $k = 1, 2, \dots, d-1$, and, in addition, the projections $p_1, p_2, \dots, p_d$ are mutually orthogonal in $C([0, 1], F_b)$, and $q_1, q_2, \dots, q_{d-1}$ mutually orthogonal in F_a .

We claim that every path in $\mathcal{G}$ of length $\leq k-1$ can be realized when $\min_i a(i) \geq k$. To prove this by induction in k , observe that a path of length 1 has the form $(i_1, *_1) \xrightarrow{j} (i_2, *_2)$ which is trivial to realize when $a(j) \geq \min_i a(i) \geq 2$. Assume then that the assertion is true for some $k \geq 2$, that $\min_i a(i) \geq k+1$ and consider a path

$$c = (i_1, *_1) \xrightarrow{j_1} (i_2, *_2) \xrightarrow{j_2} \dots \xrightarrow{j_k} (i_{k+1}, *_{k+1})$$

of length k . By induction hypothesis there are projections such that $p_1 \xrightarrow{q_1} p_2 \xrightarrow{q_2} p_3 \xrightarrow{q_3} \dots \xrightarrow{q_{k-1}} p_k$ realizes $(i_1, *_1) \xrightarrow{j_1} (i_2, *_2) \xrightarrow{j_2} \dots \xrightarrow{j_{k-1}} (i_k, *_{k+1})$. Consider first the case where $*_k = \rightarrow$ and $*_{k+1} = \rightarrow$. Since $a(j_k) \geq k+1$ there is a projection $q_k \in M_{a(j_k)}$ which is orthogonal in F_a to each $q_i, i \leq k-1$. It follows that $\varphi_{i_k j_k}^U(q_k)$ is orthogonal in F_b to $p_i(1)$ for all $i \leq k-1$, with the possible exception of $p_1(1)$ when $*_1 = \leftarrow$ and $i_1 = i_k$. When this happens we change $p_1(t)$ for t in a neighborhood of 1 to arrange that $\varphi_{i_k j_k}^U(q_k) \perp p_1(1)$, while keeping $p_1, p_2, \dots, p_k$ orthogonal. This is possible because $b(i_k) \geq \min_j a(j) \geq k+1$. So we can assume, in all cases, that $\varphi_{i_k j_k}^U(q_k)$ is orthogonal to $p_i(1)$ for all $i \leq k-1$. There is then a rank 1 projection $p'_k \in C([0, 1], M_{b(i_k)})$ such that $p'_k(t) = p_k(t), t \leq \frac{1}{2}$, $p'_k \perp p_i$ in $C([0, 1], F_b)$ for all $i \leq k-1$, and $p'_k(1) = \varphi_{i_k j_k}^U(q_k)$. Exchanging p'_k for p_k we may thus assume that $p_k(1) = \varphi_{i_k j_k}^U(q_k)$.

Note that $\varphi_{i_{k+1} j_k}^I(q_k)$ is orthogonal to $p_i(0)$ in F_b for $i \leq k$, with the possible exception of $p_1(0)$ when $i_1 = i_{k+1}$ and $*_1 = \rightarrow$. When this happens we change $p_1(t)$ for t in a neighborhood of 0 to arrange that $\varphi_{i_{k+1} j_k}^I(q_k) \perp p_1(0)$, while keeping $p_1, p_2, \dots, p_k$ orthogonal. This is possible because $b(i_{k+1}) \geq \min_j a(j) \geq k+1$. So we can assume, in all cases, that $\varphi_{i_{k+1} j_k}^I(q_k)$ is orthogonal to $p_i(0)$ for all $i \leq k$. Using once more that $b(i_{k+1}) \geq k+1$ we can find a projection p_{k+1} in $C([0, 1], M_{b(i_{k+1})})$ which is orthogonal in $C([0, 1], F_b)$ to $p_i, i \leq k$, and satisfies that $p_{k+1}(0) = \varphi_{i_{k+1} j_k}^I(q_k)$. Then $p_1 \xrightarrow{q_1} p_2 \xrightarrow{q_2} p_3 \xrightarrow{q_3} \dots \xrightarrow{q_{k-1}} p_k \xrightarrow{q_k} p_{k+1}$ realizes c . Since exactly the same proof works in the other three case, $*_k = \rightarrow$ and $*_{k+1} = \leftarrow$, $*_k = \leftarrow$ and $*_{k+1} = \rightarrow$, and finally $*_k = \leftarrow$ and $*_{k+1} = \leftarrow$, we have completed the induction step.

Note that it follows from conditions (3.3) and (3.4) that $\mathcal{G}$ has no sinks or sources. There is therefore a loop

$$c = (i_1, *_1) \xrightarrow{j_1} (i_2, *_2) \xrightarrow{j_2} \dots \xrightarrow{j_{d-1}} (i_d, *_{d-1}) \xrightarrow{j_d} (i_1, *_1),$$

in $\mathcal{G}$ whose length ($= d$) does not exceed $2n$. Set $(i_{d+1}, *_{d+1}) = (i_1, *_1)$. Since we assume that $a(i) \geq 2n + 1$ for all i , it follows from the preceding that there are projections such that $p_1 \xrightarrow{q_1} p_2 \xrightarrow{q_2} p_3 \xrightarrow{q_3} \dots \xrightarrow{q_d} p_{d+1}$ realizes c . Assume $*_1 = \leftarrow$. Change p_1 to p'_1 such that $p'_1(t) = p_1(t)$, $t \leq \frac{1}{2}$, and $p'_1(1) = \varphi_{i_1 j_d}^U(q_d)$. Since $\varphi_{i_1 j_d}^U(q_d)$ is orthogonal to $p_j(1)$, $2 \leq j \leq d$, in F_b and p_1 to p_j , $2 \leq j \leq d$, in $C([0, 1], F_b)$, we can do this such that the elements of $\{p'_1, p_2, \dots, p_d\}$ are mutually orthogonal in $C([0, 1], F_b)$. Then $(\sum_{j=1}^d q_j, p'_1 + \sum_{j=2}^d p_j)$ is a non-zero projection in $A(a, b, I, U)$. This completes the proof because the case $*_1 = \rightarrow$ is completely analogous. $\square$

Note that there is an extension

$$0 \longrightarrow C_0(0, 1) \otimes F_b \longrightarrow A(a, b, I, U) \longrightarrow F_a \longrightarrow 0$$

which makes it easy to calculate the K -theory groups of a building block $A(a, b, I, U)$:

Lemma 3.4. $K_0(A(a, b, I, U))$ is isomorphic, as a partially ordered group, to

$$\{z = (z_1, z_2, \dots, z_m) \in \mathbb{Z}^m : (I - U)z = 0\},$$

and

$$K_1(A(a, b, I, U)) \simeq \text{coker}(I - U) = \mathbb{Z}^n / (I - U)(\mathbb{Z}^m).$$

3.2. The inductive limit decomposition. In the notation of Lemma 5.14 of [Th1], there is an open interval $J \subseteq \Gamma$, contained in some edge of Γ , with endpoints in $h^{-1}(\mathbb{V})$, such that $h^d(J) = \Gamma$ for some $d \in \mathbb{N}$, and such that $B_{\bar{h}}(\bar{\Gamma})$ is $*$ -isomorphic to $A \otimes \mathbb{K}$, where A is the inductive limit of the sequence

$$C_r^*(R^d) \subseteq C_r^*(R^{d+1}) \subseteq C_r^*(R^{d+2}) \subseteq \dots$$

Here $R^i = R_{h^i}$ is the open interval-graph relation defined by $h^i : J \rightarrow \Gamma$ and the inclusions $C_r^*(R^i) \subseteq C_r^*(R^{i+1})$ are induced by considering R^i as an open sub-relation of R^{i+1} . To give a more detailed description of the sequence (3.2) we need some terminology which we now introduce. A $*$ -homomorphism $\varphi : C([0, 1], M_n) \rightarrow C([0, 1], M_m)$ is *regular* when there are continuous functions $g_1, g_2, \dots, g_k : [0, 1] \rightarrow [0, 1]$ such that

$$\varphi(f) = \text{diag}(f \circ g_1, f \circ g_2, \dots, f \circ g_k, 0, \dots, 0).$$

Of course, this requires that $kn \leq m$. φ is unital when $kn = m$. The functions $g_1, \dots, g_k$ will be called *the characteristic functions* of φ . A $*$ -homomorphism $\chi : \bigoplus_{i=1}^N C([0, 1], M_{n(i)}) \rightarrow \bigoplus_{i=1}^M C([0, 1], M_{m(i)})$ is *regular* when the $*$ -homomorphisms $\chi_{ij} : C([0, 1], M_{n(j)}) \rightarrow C([0, 1], M_{m(i)})$, $i = 1, 2, \dots, M$, $j = 1, 2, \dots, N$, which it defines are all regular. The union of the characteristic functions of the χ_{ij} 's will be called *the characteristic functions* of χ . A $*$ -homomorphism $\chi : \bigoplus_{j=1}^N M_{n(j)} \rightarrow \bigoplus_{j=1}^M M_{m(j)}$ will be said to be *non-increasing in rank* when its multiplicity matrix $A^\chi = (A_{ij}^\chi)$ is given by a map $\iota : \{1, 2, \dots, N\} \rightarrow \{1, 2, \dots, M\}$ such that

$$A_{ij}^\chi = \begin{cases} 1 & \text{when } \iota(j) = i \\ 0 & \text{otherwise.} \end{cases}$$

Let $A(a, b, I, U)$ and $A(a', b', I', U')$ be building blocks. A $*$ -homomorphism

$$\psi : A(a, b, I, U) \rightarrow A(a', b', I', U') \quad (3.5)$$

is *regular* when there are $*$ -homomorphisms $\mu : C([0, 1], F_b) \rightarrow F_{b'}$, $\chi : F_a \rightarrow F_{a'}$ and a regular $*$ -homomorphism $\varphi : C([0, 1], F_b) \rightarrow C([0, 1], F_{b'})$ such that

- 1) $\mu(f)\chi(x) = \chi(x)\mu(f) = 0$, $f \in C([0, 1], F_b)$, $x \in F_a$,
- 2) $\psi(x, f) = (\chi(x) + \mu(f), \varphi(f))$, $(x, f) \in A(a, b, I, U)$, and
- 3) χ is non-increasing in rank.

We will refer to the $*$ -homomorphism μ as *the skew map* of ψ . Note that the composition of two regular $*$ -homomorphisms is again a regular $*$ -homomorphism.

Let F be a finite-dimensional C^* -algebra. We define the *spectral variation* $\text{var } f$ of an element $f \in C([0, 1], F)$ to be the number

$$\text{var } f = \inf_U \sup_{s, t \in [0, 1]} \|U(s)f(s)U(s)^* - f(t)\|,$$

where we take the infimum over all unitaries U in $C([0, 1], F)$. The *spectral variation* $\text{var } z$ of an element $z = (x, f) \in A(a, b, I, U)$ of a building block is then defined to be the number $\text{var } z = \text{var } f$.

Lemma 3.5. *The heteroclinic algebra $B_h^-(\bar{\Gamma})$ is $*$ -isomorphic to the inductive limit of a sequence of building blocks*

$$A'_1 \xrightarrow{\pi_1} A'_2 \xrightarrow{\pi_2} A'_3 \xrightarrow{\pi_3} \dots \quad (3.6)$$

and injective regular $*$ -homomorphisms with the property that

$$\lim_{k \rightarrow \infty} \text{var} [\pi_k \circ \pi_{k-1} \circ \pi_{k-2} \circ \dots \circ \pi_i(a)] = 0 \quad (3.7)$$

for all $a \in A'_i$ and all $i \in \mathbb{N}$.

Proof. For each $i \geq d$ we let A'_i be the building block obtained from $C_r^*(R_{h^i})$ by use of Lemma 3.2. Define $\chi : \mathbb{A}_{h^i} \rightarrow \mathbb{A}_{h^{i+1}}$ such that $\chi(e_{x,y}) = e_{x,y}$ when $x, y \in h^{-i}(\mathbb{V})$ and $p_x = p_y$. This is well-defined because h maps h^i -passages to h^{i+1} -passages by the Markov condition γ) and the non folding condition β). Define $\mu : C([0, 1], \mathbb{B}_{h^i}) \rightarrow \mathbb{A}_{h^{i+1}}$ such that

$$\mu(f \otimes e_{I,J}) = \sum_{(x,y) \in N_{I,J}} f(\psi_{h^i(\bar{I})} \circ h^i(x)) e_{x,y}$$

when $f \in C[0, 1]$ and $I, J \in \mathcal{I}_{h^i}$ are such that $h^i(I) = h^i(J)$, and we sum over the set

$$N_{I,J} = \{(x, y) \in h^{-i-1}(\mathbb{V})^2 : x \in I, y \in J, h^i(x) = h^i(y)\}.$$

Define a $*$ -homomorphism $\varphi : C([0, 1], \mathbb{B}_{h^i}) \rightarrow C([0, 1], \mathbb{B}_{h^{i+1}})$ such that

$$\varphi(f \otimes e_{I,J}) = \sum_{(I_1, J_1) \in M_{I,J}} f \circ \psi_{h^i(\bar{I})} \circ (h|_{h^i(\bar{I}) \cap h^{-1}(h^{i+1}(\bar{I}_1))})^{-1} \circ \psi_{h^{i+1}(\bar{I}_1)}^{-1} \otimes e_{I_1, J_1},$$

for $f \in C[0, 1]$ and $I, J \in \mathcal{I}_{h^i}$ with $h^i(\bar{I}) = h^i(\bar{J})$, where we sum over the set

$$M_{I,J} = \{(I_1, J_1) \in \mathcal{I}_{h^{i+1}}^2 : I_1 \subseteq I, J_1 \subseteq J, h^{i+1}(\bar{I}_1) = h^{i+1}(\bar{J}_1)\}.$$

Then χ, μ, φ define a $*$ -homomorphism $\psi : \mathbb{D}_{h^i} \rightarrow \mathbb{D}_{h^{i+1}}$ such that $\psi(x, f) = (\chi(x) + \mu(f), \varphi(f))$. Note that the diagram

$$\begin{array}{ccc} \mathbb{D}_{h^i} & \xrightarrow{\psi} & \mathbb{D}_{h^{i+1}} \\ \Phi_{h^i} \uparrow & & \uparrow \Phi_{h^{i+1}} \\ C_r^*(R_{h^i}) & \subseteq & C_r^*(R_{h^{i+1}}) \end{array}$$

commutes. Set $\pi_i = \Psi_{h^{i+1}} \circ \psi \circ \Psi_{h^i}^{-1}$. Then π_i is a regular $*$ -homomorphism and the diagram

$$\begin{array}{ccc} A'_i & \xrightarrow{\pi_i} & A'_{i+1} \\ \Psi_{h^i} \circ \Phi_{h^i} \uparrow & & \uparrow \Psi_{h^{i+1}} \circ \Phi_{h^{i+1}} \\ C_r^*(R_{h^i}) & \subseteq & C_r^*(R_{h^{i+1}}) \end{array}$$

commutes. The characteristic functions of π_i are of the form

$$\psi_e \circ (h|_{\overline{I_j}})^{-1} \circ \psi_{e'}^{-1} \quad (3.8)$$

where e, e' are edges of Γ and $I_1, I_2, \dots, I_k$ are the intervals in $e \cap h^{-1}(e')$ such that $h : \overline{I_j} \rightarrow e'$ are homeomorphisms. It is then clear that the expansion condition α) on h ensures that $\lim_{k \rightarrow \infty} \text{var} [\pi_k \circ \pi_{k-1} \circ \pi_{k-2} \circ \dots \circ \pi_i(a)] = 0$ for all $a \in A'_i$ and all i . $\square$

Let H_{ij} be the number of times the edge e_i is covered by the edge e_j under h , i.e. for arbitrary $x \in \text{int } e_i$,

$$H_{ij} = \#h^{-1}(x) \cap e_j. \quad (3.9)$$

It follows from the mixing condition ϵ) that H is mixing, i.e. for large enough k the matrix H^k has no zero entries. As seen in the proof of Lemma 3.5 the sequence (3.6) is stationary in the sense that the characteristic functions of the connecting $*$ -homomorphisms, the π_i 's, are the same. They are given by continuous functions $\chi_{ij}^k : [0, 1] \rightarrow [0, 1]$, $i, j = 1, 2, \dots, n$, $k = 1, 2, \dots, H_{ij}$, where n is the number of edges in Γ , and H_{ij} is the number from (3.9). The characteristic functions have the following properties:

- n1) The characteristic functions are injective.
- n2) $\bigcup_{j,k} \chi_{ij}^k([0, 1]) = [0, 1]$ for all $i = 1, 2, \dots, n$.
- n3) When $(j, k) \neq (j', k')$ the intersection $\chi_{ij}^k([0, 1]) \cap \chi_{ij'}^{k'}([0, 1])$ consists of at most one point.
- n4) For each i, j the points $\chi_{ij}^k(0)$, $k = 1, 2, \dots, H_{ij}$, are different.

If there is a pair i, j such that there are both increasing and decreasing functions among χ_{ij}^k , $k = 1, 2, \dots, H_{ij}$, we can use the fact that H is mixing and telescope the sequence (3.6) to arrange that this is the case for all i, j . If instead it holds that for all i, j the functions χ_{ij}^k , $k = 1, 2, \dots, H_{ij}$, are either all increasing or all decreasing we telescope first (3.6) by removing every second level to make sure that, in addition, the functions χ_{11}^k , $k = 1, 2, \dots, H_{11}$, all are increasing. If this is not possible we are back in the first case so assume that χ_{11}^k , $k = 1, 2, \dots, H_{11}$, are all increasing. When j is such that the functions χ_{1j}^k , $k = 1, 2, \dots, H_{1j}$, are all decreasing we can arrange by telescoping that $\{\chi_{ij}^k\}$ contains both increasing and decreasing functions *unless* χ_{j1}^k , $k = 1, 2, \dots, H_{j1}$, are

also all decreasing. In this case we change the orientation of e_j . After this is done for each such j we telescope the sequence again by removing every second level. There is then, for each i, j , at least one k such that χ_{ij}^k is increasing. So unless we are back in the first case, all the characteristic functions must be increasing. In short, we can arrange by telescoping that

- n5) the characteristic functions are either all increasing or $\{\chi_{ij}^k : k = 1, 2, \dots, H_{ij}\}$ contains both increasing and decreasing functions for all i, j .

For convenience we shall substitute h with a power h^m of h in order to ensure further convenient properties of the connecting maps of (3.6). Such a substitution is justified by the following lemma.

Lemma 3.6. *For every $m \in \mathbb{N}$ the map $h^m : \Gamma \rightarrow \Gamma$ is a pre-solenoid, $B_{\overline{h^m}}(\overline{\Gamma}) \simeq B_{\overline{h}}(\overline{\Gamma})$ and $B_{\overline{h^m}^{-1}}(\overline{\Gamma}) \simeq B_{\overline{h}^{-1}}(\overline{\Gamma})$.*

Proof. The first statement is straightforward to check, and the others follow from the observation that $(\overline{\Gamma}, \overline{h^m})$ is conjugate to $(\overline{\Gamma}, \overline{h})$, combined with the observation that the heteroclinic algebra of a homeomorphism is $*$ -isomorphic to the heteroclinic algebra of any of its positive powers. $\square$

It follows from the mixing condition δ) that there is a $m \in \mathbb{N}$ such that $h^m(e) = \Gamma$ for every edge e in Γ . Hence, for a study of the heteroclinic algebra, we may assume, by Lemma 3.6, that

$$h(e) = \Gamma, \quad e \in \mathbb{E}. \quad (3.10)$$

In fact we are going to move to higher powers of h in order to obtain other properties. Note that this will not violate (3.10). Let $\mathcal{P}_i$ denote the h^i -passages. It follows from (3.10) that $\mathcal{P}_i \subseteq \mathcal{P}_{i+1}$. Since there are only finitely many passages in Γ it follows that there is an m so big that $\mathcal{P}_{i+m} = \mathcal{P}_m$ for all $i \geq 0$. A similar argument shows that if m is big enough the set of passages contained in $h^i(\text{int } e)$ is the same for each $i \geq m$ and each $e \in \mathbb{E}$. Then (3.10) shows that this 'stabilized' set of passages is independent of e . Hence by substituting h with h^m for some sufficiently large m we can arrange that there is a set Q of passages in Γ such that

$$Q \text{ is the set of } h^i\text{-passages for all } i \in \mathbb{N}, \quad (3.11)$$

and

$$Q \text{ is the set of passages contained in } h(\text{int } e) \text{ for all } e \in \mathbb{E}. \quad (3.12)$$

By using the properties (3.10), (3.11) and (3.12) in the proof of Lemma 3.5 we can regularize the resulting sequence of C^* -algebras further. To describe the property of the connecting maps which we obtain from (3.12), consider a regular $*$ -homomorphism as in (3.5) with skew map μ . We say that ψ is *full* when the composition

$$C_0(0, 1) \otimes M_{b(i)} \hookrightarrow C([0, 1], F_b) \xrightarrow{\mu} F_{a'} \rightarrow M_{a'(j)}$$

is non-zero for all i, j . By using the properties (3.10), (3.11) and (3.12) in the proof of Lemma 3.5, we obtain the following

Lemma 3.7. *There are natural numbers $n, m \in \mathbb{N}$ and $n \times m$ $\{0, 1\}$ -matrices I, U and a set $\chi_{ij}^k : [0, 1] \rightarrow [0, 1]$, $k = 1, 2, \dots, H_{ij}$, $i, j = 1, 2, \dots, n$, of continuous functions such*

that $H_{ij} \geq 1$ for all i, j , and sequences $\{a_i\} \subseteq \mathbb{N}^m$, $\{b_i\} \subseteq \mathbb{N}^n$, such that $B_{\overline{h}}(\overline{\Gamma})$ is the inductive limit of a sequence

$$A(a_1, b_1, I, U) \xrightarrow{\pi_1} A(a_2, b_2, I, U) \xrightarrow{\pi_2} A(a_3, b_3, I, U) \xrightarrow{\pi_3} \dots \quad (3.13)$$

such that (3.7) holds and each π_i is a full regular $*$ -homomorphism with $\{\chi_{ij}^k\}$ as characteristic functions.

Lemma 3.8. *Let $\varphi : A \rightarrow B$ be a regular $*$ -homomorphism between building blocks A and B . Let $e \in A$ be a projection. There are unital building blocks, A' and B' , a unital regular $*$ -homomorphism $\varphi' : A' \rightarrow B'$ and $*$ -isomorphisms $eAe \rightarrow A'$ and $\varphi(e)B\varphi(e) \rightarrow B'$ such that*

$$\begin{array}{ccc} eAe & \longrightarrow & A' \\ \varphi \downarrow & & \downarrow \varphi' \\ \varphi(e)B\varphi(e) & \longrightarrow & B' \end{array} \quad (3.14)$$

commutes.

Proof. Left to the reader. □

In the following we let $\mathbb{K}$ denote the C^* -algebra of compact operators on an infinite-dimensional separable Hilbert space.

Proposition 3.9. *There are natural numbers $n, m \in \mathbb{N}$ and $n \times m$ $\{0, 1\}$ -matrices I, U and a set $\chi_{ij}^k : [0, 1] \rightarrow [0, 1]$, $k = 1, 2, \dots, H_{ij}$, $i, j = 1, 2, \dots, n$, of continuous functions such that $H_{ij} \geq 1$ for all i, j , and sequences $\{a_i\} \subseteq \mathbb{N}^m$, $\{b_i\} \subseteq \mathbb{N}^n$, such that $B_{\overline{h}}(\overline{\Gamma})$ is $*$ -isomorphic to $A \otimes \mathbb{K}$ where A is the inductive limit of a sequence*

$$A(a_1, b_1, I, U) \xrightarrow{\pi_1} A(a_2, b_2, I, U) \xrightarrow{\pi_2} A(a_3, b_3, I, U) \xrightarrow{\pi_3} \dots \quad (3.15)$$

of unital building blocks such that (3.7) holds and each π_i is a unital full regular $*$ -homomorphism with $\{\chi_{ij}^k\}$ as the characteristic functions.

Proof. Let n, m, U, I, H and $\{\chi_{ij}^k\}$ be as in Lemma 3.7. The mixing condition combined with the fullness of the connecting maps in (3.13) guarantees that

$$\lim_{i \rightarrow \infty} \min_k a_i(k) = \infty$$

and it follows therefore from Lemma 3.3 that the building blocks in (3.13) contain projections, at least from a certain stage. The fullness of the connecting maps ensure that these projections become full projections in the sense of [Br]. $B_{\overline{h}}(\overline{\Gamma})$ is simple and stable by Lemma 5.10 of [Th1] so it follows from [Br] that $B_{\overline{h}}(\overline{\Gamma}) \simeq pB_{\overline{h}}(\overline{\Gamma})p \otimes \mathbb{K}$ for any non-zero projection $p \in B_{\overline{h}}(\overline{\Gamma})$. As we have just argued we can assume that p is a full projection in the first building block occurring in (3.13). The proof is then completed by use of Lemma 3.8. □

3.3. Real rank zero and the consequences.

Lemma 3.10. *The C^* -algebra A of Proposition 3.9 has a unique trace state.*

Proof. Consider the sequence (3.15), and set $A_i = A(a_i, b_i, I, U)$. There is a $*$ -homomorphism $\iota_i : A_i \rightarrow C([0, 1], F_{b_i}) = B_i$ such that $\iota_i(x, f) = f$. Since the connecting maps are regular we get an infinite commuting diagram

$$\begin{array}{ccccccc} A_1 & \xrightarrow{\pi_1} & A_2 & \xrightarrow{\pi_2} & A_3 & \xrightarrow{\pi_3} & \cdots \\ \downarrow \iota_1 & & \downarrow \iota_2 & & \downarrow \iota_3 & & \\ B_1 & \xrightarrow{\varphi_1} & B_2 & \xrightarrow{\varphi_2} & B_3 & \xrightarrow{\varphi_3} & \cdots \end{array} \quad (3.16)$$

where the φ_i 's are regular $*$ -homomorphisms between interval algebras. It follows then from condition (3.7) that the inductive limit C^* -algebra $B = \varinjlim (B_i, \varphi_i)$ is AF. Furthermore, the sequence $K_0(B_i) \xrightarrow{\varphi_{i*}} K_0(B_{i+1})$ is stationary: $K_0(B_i) \simeq \mathbb{Z}^n$ where n is the number of edges in Γ and φ_{i*} is given by the matrix H , cf. (3.9). Since H is mixing it follows that B is a unital simple AF-algebra with a unique trace state, cf. Theorem 6.1 of [Ef].

To complete the proof we need some notation. When D is a unital C^* -algebra we denote by $T(D)$ its tracial state space and by $\text{Aff } T(D)$ the order unit space of real-valued affine continuous functions on $T(D)$. When $\nu : D \rightarrow C$ is a $*$ -homomorphism between unital C^* -algebras we let $\widehat{\nu} : D_{sa} \rightarrow \text{Aff } T(C)$ be the bounded linear map from the self adjoint part of D to $\text{Aff } T(C)$ given by

$$\widehat{\nu}(d)(\omega) = \omega(\nu(d)), \quad \omega \in T(C).$$

Note that

$$\|\widehat{\nu' \circ \nu}\| \leq \|\widehat{\nu'}\| \|\widehat{\nu}\| \quad (3.17)$$

when $\nu' : C \rightarrow E$ is another $*$ -homomorphism between unital C^* -algebras. Let $\pi_{k,i}(x, f) = (\chi_{k,i}(x) + \mu_{k,i}(f), \varphi_{k,i}(f))$ be the decomposition of $\pi_{k,i} = \pi_{k-1} \circ \pi_{k-2} \circ \cdots \circ \pi_i : A_i \rightarrow A_k$ arising from the fact that the π_j 's, and hence also $\pi_{k,i}$, are regular $*$ -homomorphisms. Let $z = (x, f) \in A_i$ be a self-adjoint element in the unit ball of A_i and let $\epsilon > 0$. We will show that there is a constant $\lambda \in [-1, 1]$ such that

$$\|\widehat{\pi_{\infty,i}}(z) - \lambda 1\| \leq 2\epsilon \quad (3.18)$$

in $\text{Aff } T(A)$, where $\pi_{\infty,i} : A_i \rightarrow A$ is the canonical embedding going with the inductive limit construction. The desired conclusion follows easily from (3.18) since i, z and $\epsilon > 0$ are arbitrary. Observe first that since B has a unique trace state there is a constant $\lambda \in [-1, 1]$ such that $\lim_{k \rightarrow \infty} \|\widehat{\varphi_{k,i}}(f) - \lambda 1\| = 0$. Hence $\|\widehat{\varphi_{j,i}}(f) - \lambda 1\| \leq \epsilon$ for all large j . Write $\pi_{j,i}(z) = (y, g)$ where $g = \varphi_{j,i}(f)$. Since $\varphi_{j,i}$ is unital it follows from (3.17) that

$$\|\widehat{\mu_{k,j}}(g) - \lambda \widehat{\mu_{k,j}}(1)\| = \|\widehat{\mu_{k,j} \circ \varphi_{j,i}}(f - \lambda 1)\| \leq \|\widehat{\varphi_{j,i}}(f) - \lambda 1\| \leq \epsilon \quad (3.19)$$

for all $k \geq j$. Since $\chi_{k,i}$ does not increase rank by condition 3) on a regular $*$ -homomorphism, we find that

$$\lim_{k \rightarrow \infty} \|\widehat{\chi_{k,j}}(y - \lambda 1)\| = 0. \quad (3.20)$$

By combining (3.19) and (3.20) we find that

$$\|\widehat{\chi_{k,j}}(y) + \widehat{\mu_{k,j}}(g) - \lambda 1\| = \|\widehat{\chi_{k,j}}(y - \lambda 1) + \widehat{\mu_{k,j}}(g) - \lambda \widehat{\mu_{k,j}}(1)\| \leq 2\epsilon$$

for all large k . It follows that (3.18) holds. $\square$

Lemma 3.11. $B_{\overline{h}}(\overline{\Gamma})$ has real rank zero.

Proof. By [BP] we must show that the C^* -algebra A of Proposition 3.9 has real rank zero. Since A is a simple unital recursively subhomogenous C^* -algebra in the sense of [Ph3] and since A only has one trace state by Lemma 3.10 it suffices by Theorem 4.2 of [Ph3] to show that there are non-zero projections in A of arbitrary small trace. For this purpose observe that the projection p constructed in the proof of Lemma 3.3 satisfies the inequality

$$\tau(p) \leq \frac{2n+1}{\min_k a(k)}$$

for all $\tau \in T(A(a, b, I, U))$. As we have already used above, the mixing condition and the fullness of the connecting $*$ -homomorphisms in (3.15) imply that the number $\min_k a(k)$ becomes arbitrarily large in the building blocks of (3.15). It follows that A has non-zero projections of arbitrarily small trace. $\square$

Remark 3.12. My original proof of Lemma 3.11 was more direct, but also more complicated. I am grateful to N.C. Phillips for his suggestion to use the result from [Ph3] to shorten the proof. $\square$

We know now that $B_{\overline{h}}(\overline{\Gamma}) \simeq A \otimes \mathbb{K}$ where A

- is the inductive limit of a sequence of unital building blocks with unital connecting maps,
- has real rank zero, and
- has a unique trace state.

This allows us to use the work of H. Lin on simple C^* -algebras of tracial rank 0.

Lemma 3.13. A has tracial rank zero in the sense of H. Lin, cf. [Lin2].

Proof. By Proposition 5.4 of [Lin3] $T(A)$ is approximately AC. It is easily shown that the invertibles are dense in a unital building block and it follows that the same is true in A , i.e. A has stable rank one. Furthermore, it follows from Lemma 3.4 that $K_0(A)$ is unperforated. Hence Theorem 4.15 of [Lin3] shows that A has tracial rank zero. $\square$

By an *AH-algebra with no dimension-growth* we mean in the following the inductive limit of a sequence of C^* -algebras of the form

$$p_1 C(X_1, M_{k_1}) p_1 \rightarrow p_2 C(X_2, M_{k_2}) p_2 \rightarrow p_3 C(X_3, M_{k_3}) p_3 \rightarrow \cdots, \quad (3.21)$$

where each X_i is a compact metric space, p_i is a projection in $C(X_i, M_{k_i})$ and $\sup_i \dim X_i < \infty$. In the case where each of the X_i 's is a disjoint union of circles the resulting algebra is called an *AT-algebra*.

Theorem 3.14. Let $B_{\overline{h}}(\overline{\Gamma})$ be the heteroclinic algebra of the 1-solenoid $(\overline{\Gamma}, \overline{h})$. Then $B_{\overline{h}}(\overline{\Gamma}) \simeq A \otimes \mathbb{K}$ where A is a unital simple AH-algebra with real rank zero, no dimension growth and a unique trace.

Proof. As pointed out by N.C. Phillips in Proposition 4.7 of [Ph2] this follows from the work of Elliott, Gong and Lin, cf. [Lin3] and [EG]. $\square$

In particular, it follows now, thanks to the work of Elliott and Gong, [EG], or of Elliott, Gong and Li, [EGL], that $B_{\bar{h}}(\bar{\Gamma})$ is classified by K-theory. In fact, since $K_0(B_{\bar{h}}(\bar{\Gamma}))$ is torsion free it follows from the concluding remark in [EGL] that $B_{\bar{h}}(\bar{\Gamma})$ can also be realized by the inductive limit of a sequence of direct sums of circle algebras and matrix algebras over so-called dimension drop algebras, i.e. algebras of the form

$$\{f \in C([0, 1], M_k) : f(0) \in \mathbb{C}, f(1) \in \mathbb{C}\}.$$

Thus $B_{\bar{h}}(\bar{\Gamma})$ is actually in the class of real rank zero algebras classified in Elliotts breakthrough paper, [Ell2]. Furthermore, since the dimension drop algebras are only needed in the presence of torsion in K_1 , it will follow from results below that the dimension drop algebras can be omitted if and only if $(\bar{\Gamma}, \bar{h})$ is orientable.

3.4. On the K-theory of $B_{\bar{h}}(\bar{\Gamma})$. With the detailed information on the connecting maps at hand it is easy to obtain the following description of the K-theory of $B_{\bar{h}}(\bar{\Gamma})$ from the 'stationary' sequence of Lemma 3.9.

Proposition 3.15. *The partially ordered group $K_0(B_{\bar{h}}(\bar{\Gamma}))$ is the inductive limit of the sequence*

$$H \xrightarrow{\chi_* + \mu_* \circ I} H \xrightarrow{\chi_* + \mu_* \circ I} H \xrightarrow{\chi_* + \mu_* \circ I} \dots,$$

where $H = \{x \in \mathbb{Z}^m : Ix = Ux\}$, and

$$K_1(B_{\bar{h}}(\bar{\Gamma})) \simeq \text{coker}(I - U) = \mathbb{Z}^n / (I - U)(\mathbb{Z}^m).$$

Proof. Set $C_k = A(a_k, b_k, I, U)$ and identify $K_0(F_{a_k})$ with $\mathbb{Z}^m$ and $K_0(F_{b_k}) = K_1(SF_{b_k})$ with $\mathbb{Z}^n$. The conclusion regarding K_0 follows from Lemma 3.9 and Lemma 3.4 by observing that

$$\begin{array}{ccccc} K_0(C_k) & \longrightarrow & \mathbb{Z}^m & \xrightarrow{I-U} & \mathbb{Z}^n \\ \pi_{k*} \downarrow & & \chi_* + \mu_* \circ I \downarrow & & \\ K_0(C_{k+1}) & \longrightarrow & \mathbb{Z}^m & \xrightarrow{I-U} & \mathbb{Z}^n \end{array}$$

commutes. Similarly, by using the properties n1)-n5) of the characteristic functions it follows that also

$$\begin{array}{ccccc} \mathbb{Z}^m & \xrightarrow{I-U} & \mathbb{Z}^n & \longrightarrow & K_1(C_k) \\ & & \parallel & & \downarrow \pi_{k*} \\ \mathbb{Z}^m & \xrightarrow{I-U} & \mathbb{Z}^n & \longrightarrow & K_1(C_{k+1}) \end{array}$$

commutes. This implies the statement concerning K_1 . $\square$

Note that $K_0(B_{\bar{h}}(\bar{\Gamma}))$ is a stationary partially ordered group, but in a slightly more general sense than the usual because H is not (obviously) a simplicial group. For this reason we must resort to Theorem 3.14 in order to conclude that $K_0(B_{\bar{h}}(\bar{\Gamma}))$ is a dimension group. In the following we focus on $K_1(B_{\bar{h}}(\bar{\Gamma}))$.

Let $A = (a_{ij})$ be an $n \times m$ matrix. Let G_A be the unoriented graph with vertex set $\{1, 2, \dots, n\}$ and an edge between i and j if and only if $i \neq j$ and there is a $k \in$

$\{1, 2, \dots, m\}$ such that $a_{ik} \neq 0$ and $a_{jk} \neq 0$. We say that A is *irreducible* when G_A is connected.

Lemma 3.16. *Let $n, m \in \mathbb{N}, n \geq 2$. Let A be an irreducible $n \times m$ integer matrix with the property that each non-zero column has exactly two non-zero entries from the set $\{1, -1\}$ or exactly one non-zero entry from the set $\{2, -2\}$.*

a) *If A contains an entry from $\{2, -2\}$ it follows that the cokernel of $A : \mathbb{Z}^m \rightarrow \mathbb{Z}^n$ is isomorphic to $\mathbb{Z}_2$.*

b) *If each non-zero column of A has exactly two non-zero entries, 1 and -1 , it follows that the cokernel of $A : \mathbb{Z}^m \rightarrow \mathbb{Z}^n$ is isomorphic to $\mathbb{Z}$.*

Proof. a) We will prove by induction in n that there is a matrix $H \in \text{Gl}_n(\mathbb{Z})$ such that $HA(\mathbb{Z}^m) = 2\mathbb{Z} \oplus \mathbb{Z}^{n-1}$. It is easy to see that this holds when $n = 2$. Consider then the case $n > 2$. By permuting the rows and columns we can arrange that $a_{11} \neq 0$ and $a_{21} \neq 0$. (The possibility that each column only has one non-zero entry is ruled out by the irreducibility of A .) If a_{11} and a_{21} have the same sign we subtract the first row from the second; otherwise we add the first row to the second. Both operations correspond to the multiplication by an element of $\text{Gl}_n(\mathbb{Z})$. The resulting $n \times m$ -matrix A' has the form

$$A' = \begin{pmatrix} \pm 1 & * \\ 0 & B \end{pmatrix}, \quad (3.22)$$

where $*$ is an $1 \times (m-1)$ matrix with entries from $\{-2, -1, 0, 1, 2\}$, the 0 is $(n-1) \times 1$ and B is an irreducible $(n-1) \times (m-1)$ integer matrix for which we can use the induction hypothesis. In particular B contains an entry from $\{2, -2\}$. Let $H_0 \in \text{Gl}_{n-1}(\mathbb{Z})$ be such that $H_0 B(\mathbb{Z}^{m-1}) = 2\mathbb{Z} \oplus \mathbb{Z}^{n-2}$. Then

$$H = \begin{pmatrix} 1 & 0 \\ 0 & H_0 \end{pmatrix} \in \text{Gl}_n(\mathbb{Z})$$

has the property that $HA'(\mathbb{Z}^m) = \mathbb{Z} \oplus 2\mathbb{Z} \oplus \mathbb{Z}^{n-2}$. This completes the induction step and hence the proof.

b) is proved in the same way: Use induction to show that there is a matrix $H \in \text{Gl}_n(\mathbb{Z})$ such that $HA(\mathbb{Z}^m) = 0 \oplus \mathbb{Z}^{n-1}$. $\square$

Lemma 3.17. *Let $n, m \in \mathbb{N}, n \geq 2$. Let A be an irreducible $n \times m$ integer matrix with the property that each non-zero column has exactly two non-zero entries from the set $\{1, -1\}$. It follows that the cokernel of $A : \mathbb{Z}^m \rightarrow \mathbb{Z}^n$ is isomorphic to $\mathbb{Z}$ or to $\mathbb{Z}_2$.*

Proof. After appropriate permutations of rows and columns and a single permissible row operation we obtain a matrix A' as in (3.22) such that the matrix B is covered by a) or b) in Lemma 3.16. $\square$

Theorem 3.18. *Let $B_{\bar{h}}(\bar{\Gamma})$ be the heteroclinic algebra of the 1-solenoid $(\bar{\Gamma}, \bar{h})$. Then $K_1(B_{\bar{h}}(\bar{\Gamma})) \simeq \mathbb{Z}$ when $h : \Gamma \rightarrow \Gamma$ is oriented and $K_1(B_{\bar{h}}(\bar{\Gamma})) \simeq \mathbb{Z}_2$ when it is not.*

Proof. When h is oriented there is an orientation of the edges of Γ such that h^2 is orientation preserving and then the matrix $U - I$ will have the property described in b) of Lemma 3.16 and hence $K_1(B_{\bar{h}}(\bar{\Gamma})) \simeq \mathbb{Z}$.

Assume that $K_1(B_{\overline{h}}(\overline{\Gamma})) \simeq \mathbb{Z}$. Let Q be the passages of (3.12). Let e, f be two edges of Γ . The passages from Q which involve both e and f are of the form

$$\begin{aligned} + & : \quad \xrightarrow{e} \xrightarrow{f} \quad \text{or} \quad \xrightarrow{f} \xrightarrow{e}, \\ - & : \quad \xrightarrow{e} \xleftarrow{f} \quad \text{or} \quad \xleftarrow{e} \xrightarrow{f}. \end{aligned}$$

We use these parities to define a labeled unoriented graph H whose vertexes are the edges of Γ . Let $e, f \in \mathbb{E}$. If all passages involving e and f are of parity $+$ draw an edge between e and f with label $+$. If not, and there is in fact at least one passage involving both e and f , draw an edge between e and f with label $-$. Note that H is connected. A change in the orientation of the edges in Γ will not change the graph H - only its labels. Let $\mathbb{G}$ denote the collection of connected subgraphs H' of H with the property that there is an orientation of the edges in Γ such that all edges of H' are labeled $+$. Then $\mathbb{G}$ is not empty and is partially ordered by inclusion. Let $H_{\max}$ be a maximal element. If $H_{\max} \neq H$ there must be a loop in H of the form

$$e_1 \xrightarrow{-} e_2 \xrightarrow{+} e_3 \xrightarrow{+} e_4 \xrightarrow{+} \dots \xrightarrow{+} e_1$$

Writing down what this means for $U - I$ it is easy to see that we can obtain a matrix B from $U - I$ by permutations and addition and/or subtraction of rows such that

$$B = \begin{pmatrix} 1 & * & * & * & \dots & * \\ 0 & 1 & * & * & \dots & * \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \dots & B_0 \end{pmatrix}$$

where B_0 is an irreducible matrix for which a) Lemma 3.16 applies. It follows that $K_1(B_{\overline{h}}(\overline{\Gamma})) \simeq \mathbb{Z}_2$ which contradicts the assumption, proving that $H_{\max} = H$.

Give the edges of Γ an orientation such that all the passages, Q , are of type $+$. It follows that for all large enough i , $h^i : J \rightarrow \Gamma$ will either be orientation preserving or orientation reversing. If $h^i : J \rightarrow \Gamma$ and $h^{i+1} : J \rightarrow \Gamma$ are either both orientation preserving or both orientation reversing, $h : \Gamma \rightarrow \Gamma$ will be orientation preserving. Otherwise h must be orientation reversing. It follows that h is either positively or negatively oriented.

We have shown that (Γ, h) is oriented if and only if $K_1(B_{\overline{h}}(\overline{\Gamma})) \simeq \mathbb{Z}$. It follows then from Lemma 3.17 that $K_1(B_{\overline{h}}(\overline{\Gamma})) \simeq \mathbb{Z}_2$ when (Γ, h) is not oriented. $\square$

Theorem 3.19. *Assume that the pre-solenoid $h : \Gamma \rightarrow \Gamma$ is oriented. It follows that the heteroclinic algebra $B_{\overline{h}}(\overline{\Gamma})$ is an AT-algebra.*

Proof. By Theorem 3.18 and Proposition 3.15 the K-theory of $B_{\overline{h}}(\overline{\Gamma})$ is torsion-free. Therefore the conclusion follows from Theorem 3.14 and Theorem [EG]. $\square$

We can now give the proof of Theorem 2.2:

Proof. Let (Γ, h) be a negatively orientable pre-solenoid. There is then a negatively oriented pre-solenoid (Γ', h') such that $(\overline{\Gamma}, \overline{h})$ is conjugate to $(\overline{\Gamma'}, \overline{h'})$. Then $B_{\overline{h}}(\overline{\Gamma})$ is $*$ -isomorphic to $B_{\overline{h'}}(\overline{\Gamma'})$ and hence the two algebras have the same K_1 -group. It follows therefore from Theorem 3.18 that (Γ, h) is oriented. Assume to reach a contradiction that (Γ, h) is positively oriented. Give Γ an orientation such that h is positively oriented

with respect to it, and give Γ' an orientation such that h' is negatively oriented with respect to it. Then both $h^2 : \Gamma \rightarrow \Gamma$ and $h'^2 : \Gamma' \rightarrow \Gamma'$ preserve the orientation and hence $\bar{\Gamma}$ and $\bar{\Gamma}'$ are both oriented matchbox manifolds in the sense of [AO]. Since (Γ, h) and (Γ', h') are conjugate there is a homeomorphism $\mu : \bar{\Gamma} \rightarrow \bar{\Gamma}'$ such that

$$\bar{h}' \circ \mu = \mu \circ \bar{h}. \quad (3.23)$$

The 'matchboxes' in $\bar{\Gamma}$ and $\bar{\Gamma}'$, in the sense of [AO], are all the product of $[-1, 1]$ and the Cantor set. Since there are no isolated points in the Cantor set it is easy to see that μ must either preserve or invert the orientation. None of the two possibilities are consistent with (3.23) since $\bar{h}'$ inverts the orientation of Γ' while $\bar{h}$ preserves the orientation of $\bar{\Gamma}$. This contradiction shows that (Γ, h) must be negatively oriented.

The case where (Γ, h) is a positively orientable pre-solenoid is handled in exactly the same way. $\square$

4. THE HETEROCLINIC ALGEBRA OF THE INVERSE OF A 1-SOLENOID

We seek here to clarify the structure of the heteroclinic algebra of the inverse of $\bar{h}$, i.e. we seek to describe $B_{\bar{h}^{-1}}(\bar{\Gamma})$. Since $(\bar{\Gamma}, \bar{h})$ is a Smale space, $B_{\bar{h}^{-1}}(\bar{\Gamma})$ is the same algebra as the stabilization of the 'unstable algebra' of Putnam, cf. [Pu] and Section 4.4 of [Th1]. However, the technicalities required to show that a 1-solenoid has a Smale space structure are quite involved and tedious, and do not seem to exist in the literature. For this reason we continue to ignore this additional structure since it is not really needed. Only in Section 5 below will it be necessary to appeal to the Smale space structure.

4.1. Simplicity and stability. In this section we show that $B_{\bar{h}^{-1}}(\bar{\Gamma})$ is simple and stable.

Lemma 4.1. *Let $x \in W_{\bar{\Gamma}, \bar{h}^{-1}}$. The sets*

$$U(x, n) = \{y \in \bar{\Gamma} : y_n = x_n\}, \quad n \in \mathbb{N},$$

form a neighborhood base of x for the Wagoner topology of $W_{\bar{\Gamma}, \bar{h}^{-1}}$.

Proof. Consider the $\kappa > 0$ and the $l \in \mathbb{N}$ from Lemma 2.9 of [Y1]. When $y \in U(x, n)$, $z \in \bar{\Gamma}$ and $D(\bar{h}^m(z), \bar{h}^m(y)) < \kappa$, $m \geq -l - n$, it follows that $d(h^k(z_{l+n}), h^k(y_{l+n})) < \kappa$ for all $k \in \mathbb{N}$. By Lemma 2.9 of [Y1] this implies that $y_n = h^l(y_{l+n}) = h^l(z_{l+n}) = z_n$, proving that

$$\{z \in \bar{\Gamma} : D(\bar{h}^m(z), \bar{h}^m(y)) < \kappa, \quad m \geq -l - n\} \subseteq U(x, n).$$

By the definition of the Wagoner topology of $W_{\bar{\Gamma}, \bar{h}^{-1}}$ this implies that $U(x, n)$ is open in $W_{\bar{\Gamma}, \bar{h}^{-1}}$, cf. Lemma 4.6 [Th1]. To prove that the $U(x, n)$'s form a base consider an element

$$\Omega = \{y \in \bar{\Gamma} : D(\bar{h}^m(y), \bar{h}^m(x)) < \epsilon, \quad m \geq k\}$$

of the base described in Lemma 4.6 of [Th1]. Choose $L \in \mathbb{N}$ such that

$$\sum_{j=L+1}^{\infty} 2^{-j} \sup_{a, b \in \Gamma} d(a, b) < \epsilon.$$

Then $U(x, |k| + L) \subseteq \Omega$. □

Lemma 4.2. *Let $x, y \in \bar{\Gamma}$ be post-periodic for $\bar{h}^{-1}$. Assume that there is an $N \in \mathbb{N}$ such that*

$$\forall i \geq N \exists e_i \in \mathbb{E} : [x_i, y_i] \subseteq \text{int } e_i.$$

It follows that x is locally conjugate to y .

Proof. Let $x' \in U(x, N)$. Note that there is an edge f_{N+1} such that $x'_{N+1} \in \text{int } f_{N+1}$ and that there is a unique interval $[x'_{N+1}, y'_{N+1}] \subseteq \text{int } f_{N+1}$ such that $h([x'_{N+1}, y'_{N+1}]) = [x_N, y_N]$. Continuing recursively we obtain edges $f_i, i \geq N+1$, and intervals $[x'_i, y'_i] \subseteq \text{int } f_i$ such that $h([x'_i, y'_i]) = [x'_{i-1}, y'_{i-1}]$ for all $i \geq N+2$. We define $\chi : U(x, N) \rightarrow U(y, N)$ such that

$$\chi(x')_i = \begin{cases} y_i, & i \leq N \\ y'_i, & i \geq N+1. \end{cases}$$

Note that $\chi(x) = y$. By symmetry we can define $\mu : U(y, N) \rightarrow U(x, N)$ in a similar way and it is then clear that $\chi \circ \mu = \text{id}_{U(y, N)}$ while $\mu \circ \chi = \text{id}_{U(x, N)}$. Furthermore, it follows easily from Lemma 4.1 that both χ and μ are continuous for the Wagoner topology. Finally, it follows from the expansion condition that there are constants $C' > 0$ and $\lambda > 1$ such that $d(x'_i, \chi(x')_i) \leq C' \lambda^{-i}$ for all $x' \in U(x, N)$ and all $i \geq N$. In particular,

$$\lim_{n \rightarrow \infty} \sup_{x' \in U(x, N)} D(\bar{h}^{-n}(x'), \bar{h}^{-n}(\chi(x'))) = 0,$$

completing the proof that $(U(x, N), U(y, N), \chi)$ is a local conjugacy from x to y . □

Lemma 4.3. *Let $x, y \in \bar{\Gamma}$ be post-periodic for $\bar{h}^{-1}$. Assume that $x_N \notin \mathbb{V}$ for some N . For every $n \in \mathbb{N}$ there is an element $z \in U(y, n)$ such that x is locally conjugate to z .*

Proof. It follows from the mixing condition that there is a $k \in \mathbb{N}$ such that $h^k(\text{int } e) = \Gamma$ for all $e \in \mathbb{E}$. Choose $e \in \mathbb{E}$ such that $x_m \in \text{int } e$, where $m = \max\{n, N\} + k$. There is then an element $z_m \in \text{int } e$ such that $h^k(z_m) = y_{m-k}$. We can then define $z_i, i \geq m+1$, recursively by the requirement that $[z_i, x_i] \cap \mathbb{V} = \emptyset$ and $h(z_i) = z_{i-1}$. Set $z_j = h^{m-j}(z_m), j \leq m$. Then $z = (z_j)_{j=0}^\infty \in U(y, n)$ and it follows from Lemma 4.2 that z is locally conjugate to x . □

Remark 4.4. With future investigations in mind I want to point out that the last three lemmas, unlike the following, do not depend on the flattening condition ϵ .

Thanks to Lemma 3.6, in order to study the structure of $B_{\bar{h}^{-1}}(\bar{\Gamma})$ we may substitute h with any positive power of h . As in [Y4] we can use this together with a result of Williams, 5.2 of [Wi2], to restrict our attention to the case where Γ is the wedge of n circles, each of which is an edge in Γ , such that the common point of the circles is the only vertex in Γ . To emphasize this assumption we denote the graph by Γ_c . Let $e_i, i = 1, 2, \dots, n$, be the edges of Γ_c and let v be the unique vertex in Γ_c . By the flattening condition ϵ) and Lemma 3.6 we can also assume that there is an open neighborhood U of v in Γ_c such that

$h(U) \simeq] - 1, 1[$ and that $h(e_i) = \Gamma_c$ for all i . There are numbers $a_{ij} \in \mathbb{N}$ and disjoint intervals $I_{ij}^l, l = 1, 2, \dots, a_{ij}$, such that

$$\text{int}(e_i) \cap h^{-1}(\text{int}(e_j)) = \bigcup_{l=1}^{a_{ij}} I_{ij}^l. \quad (4.1)$$

Set $\mathcal{J} = \{I_{ij}^l : l = 1, 2, \dots, a_{ij}, i, j = 1, 2, \dots, n\}$. Since we assume that $h(U) \simeq] - 1, 1[$ for all small neighborhoods U of v we can renumber the edges of Γ_c and the intervals in $\mathcal{J}$ to arrange that for some $j \in \{1, n\}$ it holds that

$$h(U) \subseteq \overline{I_{11}^1} \cup \overline{I_{jj}^{a_{jj}}} \quad (4.2)$$

or

$$h(U) \subseteq \overline{I_{1j}^1} \cup \overline{I_{j1}^{a_{jj}}}$$

If the latter occurs and $j \neq 1$, or if $1 = j$ and $h(I_{11}^1) \supseteq I_{jj}^{a_{jj}}$, we substitute h with h^2 , as we may by Lemma 3.6, to ensure that (4.2) holds and that

$$h(\overline{I_{11}^1}) \supseteq \overline{I_{11}^1}, \quad h(\overline{I_{jj}^{a_{jj}}}) \supseteq \overline{I_{jj}^{a_{jj}}}. \quad (4.3)$$

Lemma 4.5. *Let $x, y \in \overline{\Gamma_c}$ be such that $x_i = v, y_i \neq v$ for all i , and $\lim_{i \rightarrow \infty} d(x_i, y_i) = 0$. It follows that x and y are locally conjugate in $W_{\Gamma_c, \bar{h}^{-1}}$.*

Proof. Since (4.2) and (4.3) hold there is an $N \in \mathbb{N}$ such that $y_i \in I_{11}^1$ for all $i \geq N$ or $y_i \in I_{jj}^{a_{jj}}$ for all $i \geq N$. Without loss of generality we assume that $y_i \in I_{11}^1$ for all $i \geq N$. We define $\chi : U(y, N) \rightarrow U(x, N)$ in the following way. Set $\chi(y) = x$. When $z \in U(y, N) \setminus \{y\}$, set $i_z = \max\{j \in \mathbb{N} : z_i = y_i, i \leq j\}$. Then $z_{i_z+1} \in \text{int } J_1$ for some $J_1 \in \mathcal{J}$ and there is a unique point $x'_{i_z+1} \in \overline{J_1}$ such that $h(x'_{i_z+1}) = x_{i_z} = v$ and $h([z_{i_z+1}, x'_{i_z+1}]) = [z_{i_z}, x_{i_z}] \subseteq \overline{I_{11}^1}$. We can then continue recursively: Choose $x'_{i_z+2} \in J_2 \in \mathcal{J}$ such that $h(x'_{i_z+2}) = x'_{i_z+1}$ and $h([z_{i_z+2}, x'_{i_z+2}]) = [z_{i_z+1}, x'_{i_z+1}] \subseteq \overline{J_1}$ and so on. We define

$$\chi(z)_j = \begin{cases} x_j = v, & j \leq i_z, \\ x'_j, & j \geq i_z + 1. \end{cases}$$

Since $\chi(z)_j$ only depends on z_j it follows from Lemma 4.1 that χ is continuous for the Wagoner topology. Note that the expansion condition ensures that

$$\lim_{n \rightarrow \infty} \sup_{z \in U(x, N)} D(\bar{h}^{-n}(\chi(z)), \bar{h}^{-n}(z)) = 0. \quad (4.4)$$

We define a map $\mu : U(x, N) \rightarrow U(y, N - 1)$ in a similar way: Set $\mu(x) = y$. For $z \in U(x, N) \setminus \{x\}$, set $i_z = \max\{j \in \mathbb{N} : z_i = x_i = v, i \leq j\}$. Since (4.2) holds there is a unique pair $(J'_1, y'_{i_z+1}) \in \mathcal{J} \times \Gamma_c$ such that $z_{i_z+1} \in \overline{J'_1}$, $y'_{i_z+1} \in J'_1$, $h^2(y'_{i_z+1}) = y_{i_z-1}$ and $h^2([y'_{i_z+1}, z_{i_z+1}]) = [z_{i_z-1}, y_{i_z-1}] \subseteq \overline{I_{11}^1}$. There is then a unique $J'_2 \in \mathcal{J}$ such that $z_{i_z+2} \in J'_2$ and a unique element $y'_{i_z+2} \in J'_2$ such that $h(y'_{i_z+2}) = y'_{i_z+1}$ and $h([y'_{i_z+2}, z_{i_z+2}]) =$

$[z_{i_z+1}, y'_{i_z+1}] \subseteq \overline{J'_1}$. Again we can continue and we define

$$\mu(z)_j = \begin{cases} y_j, & j \leq i_z - 1 \\ h(y'_{i_z+1}), & j = i_z, \\ y'_j, & j \geq i_z + 1. \end{cases}$$

Since $\mu(z)_j$ only depends on z_{j+1} it follows from Lemma 4.1 that μ is continuous for the Wagoner topology, and as above the expansion condition ensures that

$$\lim_{n \rightarrow \infty} \sup_{z \in U(y, N)} D(\overline{h}^{-n}(\mu(z)), \overline{h}^{-n}(z)) = 0. \quad (4.5)$$

By combining (4.4) and (4.5) we find that

$$\lim_{n \rightarrow \infty} \sup_{z \in U(y, N)} D(\overline{h}^{-n}(\mu \circ \lambda(z)), \overline{h}^{-n}(z)) = 0.$$

The expansiveness of $\overline{h}$ on $\overline{\Gamma_c}$ and the definition of the Wagoner topology therefore imply that $\mu \circ \lambda(z) = z$ for all z in a Wagoner-neighborhood of y . Since we also have that

$$\lim_{n \rightarrow \infty} \sup_{z \in U(x, N) \cap \mu^{-1}(U(y, N))} D(\overline{h}^{-n}(\lambda \circ \mu(z)), \overline{h}^{-n}(z)) = 0,$$

we conclude in the same way that $\lambda \circ \mu(z) = z$ for all z in a Wagoner-neighborhood of x . It follows that λ , suitably restricted, is a local conjugacy from y to x . $\square$

Lemma 4.6. *The heteroclinic algebra $B_{\overline{h}^{-1}}(\overline{\Gamma_c})$ of $(\overline{\Gamma_c}, \overline{h}^{-1})$ is stable and simple.*

Proof. It follows from Lemma 4.3 and Lemma 4.5 that every element of $W_{\overline{\Gamma_c}, \overline{h}^{-1}}$ is locally conjugate to an element of any given open set. It follows then from Proposition 4.7 of [Re] that $B_{\overline{h}^{-1}}(\overline{\Gamma_c})$ is simple. That $B_{\overline{h}^{-1}}(\overline{\Gamma_c})$ is stable follows also from Lemma 4.3 and Lemma 4.5 by the method used to prove Lemma 5.10 of [Th1]; specifically the two lemmas show that for any compact subset K of $W_{\overline{\Gamma_c}, \overline{h}^{-1}}$ there are periodic points and local conjugacies such that a), b), c) and d) from the proof of Lemma 5.10 of [Th1] are all satisfied, and then the results of Hjelmberg and Rørdam, [HR], imply that $B_{\overline{h}^{-1}}(\overline{\Gamma_c})$ is stable. $\square$

4.2. A free minimal action by the infinite dihedral group. We retain in this section the assumptions about (Γ_c, h) made in the previous section. In particular, Γ_c is a wedge of circles. For each $e \in \mathbb{E}$ choose homeomorphisms $\psi_{\pm}^e : [0, 1] \rightarrow e$ such that $\psi_+^e(t) = \psi_-^e(1 - t)$, $t \in [0, 1]$. We will consider e as an oriented edge with an orientation defined such that ψ_+^e is orientation-preserving and hence such that ψ_-^e is orientation-reversing. The orientations we choose are basically completely arbitrary, but for notational convenience we will ensure that

$$\overline{I_{11}^1} = \psi_-^{e_1} \left(\left[0, \frac{1}{4} \right] \right) \text{ and } \overline{I_{jj}^{a_{jj}}} = \psi_+^{e_j} \left(\left[0, \frac{1}{4} \right] \right). \quad (4.6)$$

Set

$$J = \psi_-^{e_1} \left(\left[0, \frac{1}{4} \right] \right) \cup \psi_+^{e_j} \left(\left[0, \frac{1}{4} \right] \right).$$

Then

$$h \text{ is injective on } J \text{ and } h(J) \supseteq J. \quad (4.7)$$

For each $n = 0, 1, 2, \dots$, let $\mathcal{I}^n$ denote the connected components of $\Gamma_c \setminus h^{-n}(v)$. In particular, $\mathcal{I}^0$ consists of the interiors, $\text{int } e$, of the edges e in Γ_c , and $\mathcal{I}^1 = \mathcal{J}$. Note that with C and λ the constants from the expansion condition we have the estimate

$$\text{diam } \sigma \leq KC\lambda^{-n} \quad (4.8)$$

for all $\sigma \in \mathcal{I}^n$, where $K = \text{diam } \Gamma_c$ is the diameter of Γ_c .

Let Σ denote the collection of sequences

$$\sigma = (\pm, \sigma_0, \sigma_1, \sigma_2, \sigma_3, \dots),$$

where σ_j is an interval in Γ_c with $\text{int } \sigma_j \in \mathcal{I}^j$ and $h(\sigma_{j+1}) = \sigma_j$ for all j . The first entry, $+$ or $-$, in σ will be called *the sign of σ* and will be denoted by $\text{sign}(\sigma)$. We consider Σ as a compact Hausdorff space in the product topology.

Note that for all $i \geq 1$, the set

$$\{I \in \mathcal{I}^i : v \in \bar{I} \subseteq J\}$$

consists of exactly two elements, say J_i^+ and J_i^- . We choose the superscripts such that $J_i^+ \subseteq \psi_+^{e_j}([0, \frac{1}{4}])$ and $J_i^- \subseteq \psi_-^{e_1}([0, \frac{1}{4}])$. It follows then from (4.3) that $h(J_{i+1}^\pm) = J_i^\pm$. Set $a_0^- = h(I_{11}^1) = \text{int } e_1$ and

$$a_i^- = J_i^-, \quad i \geq 1.$$

Alternatively $a_i^- \in \mathcal{I}^i$ is determined, for all i , by the requirement that $a_i^- \cap \psi_-^{e_1}([0, \frac{1}{4}])$ is a non-degenerate interval which contains v . Define an element $a^- \in \Sigma$ such that

$$a^- = (\text{sign}(a^-), a_0^-, a_1^-, a_2^-, \dots) \in \Sigma,$$

where $\text{sign}(a^-)$ is determined by the condition that

$$\lim_{t \uparrow 1} \left(h|_{a_1^-} \right)^{-1} \left(\psi_{\text{sign}(a^-)}^{a_0^-}(t) \right) = v.$$

Similarly we can define an element

$$b^- = (\text{sign}(b^-), b_0^-, b_1^-, b_2^-, \dots)$$

in Σ by the requirements that $b_0^- = h(I_{jj}^{a_{jj}}) = \text{int } e_j$,

$$b_i^- = J_i^+, \quad i \geq 1,$$

and

$$\lim_{t \uparrow 1} \left(h|_{b_1^-} \right)^{-1} \left(\psi_{\text{sign}(b^-)}^{b_1^-}(t) \right) = v.$$

It is easy to check that $a^-, b^- \in \Sigma$ have the following properties:

- Lemma 4.7.** i) $a_0^- = e_1$, $b_0^- = e_j$, $\text{sign}(a^-) = +$ and $\text{sign}(b^-) = -$.
 ii) $\lim_{t \uparrow 1} \left(h^k|_{a_k^-} \right)^{-1} (\psi_+^{e_1}(t)) = v$ and $\lim_{t \uparrow 1} \left(h^k|_{b_k^-} \right)^{-1} (\psi_-^{e_j}(t)) = v$ for all $k \in \mathbb{N}$.
 iii) If $\sigma \in \Sigma$ is an element such that $v \in \sigma_i$ for all i and

$$\lim_{t \uparrow 1} (h|_{\sigma_1})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(t) \right) = v,$$

then $\sigma \in \{a^-, b^-\}$.

Let $S : \Sigma \rightarrow \Sigma$ be the continuous involution defined such that

$$S(\pm, \sigma_0, \sigma_1, \sigma_2, \sigma_3, \dots) = (\mp, \sigma_0, \sigma_1, \sigma_2, \sigma_3, \dots).$$

Set $a^+ = S(a^-)$, $b^+ = S(b^-)$. It follows from Lemma 4.7 that a^+, b^+ are the only elements σ in Σ such that $v \in \sigma_j$ for all $j = 0, 1, 2, \dots$, and $\lim_{s \downarrow 0} (h|_{\sigma_1})^{-1} \circ \psi_{\text{sign}(\sigma)}^{\sigma_0}(s) = v$. Let $\sigma \in \Sigma \setminus \{a^-, b^-\}$. By Lemma 4.7

$$j_\sigma = \min \left\{ k \in \mathbb{N} : \lim_{t \uparrow 1} (h^k|_{\sigma_k})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(t) \right) \neq v \right\} \quad (4.9)$$

is finite. Then $\lim_{t \uparrow 1} (h^k|_{\sigma_k})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(t) \right) \neq v$ when $k \geq j_\sigma$ and there is a unique element $\sigma'_k \in \mathcal{I}^k$ such that

$$\lim_{t \uparrow 1} (h^k|_{\sigma_k})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(t) \right) = \sigma_k \cap \sigma'_k. \quad (4.10)$$

Set $\sigma' = (\text{sign}(\sigma'), \sigma'_0, \sigma'_1, \sigma'_2, \dots)$, where $\sigma'_i, i < j_\sigma$, is determined by the condition that $\sigma'_i = h^{j_\sigma - i}(\sigma'_{j_\sigma})$ and $\text{sign}(\sigma')$ is determined by the condition that

$$\lim_{t \uparrow 1} (h^i|_{\sigma_i})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(t) \right) = \lim_{s \downarrow 0} (h^i|_{\sigma'_i})^{-1} \left(\psi_{\text{sign}(\sigma')}^{\sigma'_0}(s) \right) \quad (4.11)$$

for all $i \geq 1$. We define a map $\varphi : \Sigma \rightarrow \Sigma$ such that

$$\varphi(\sigma) = \begin{cases} \sigma', & \text{when } \sigma \notin \{a^-, b^-\} \\ b^+, & \text{when } \sigma = a^- \\ a^+, & \text{when } \sigma = b^- \end{cases}$$

Lemma 4.8. φ is a homeomorphism of Σ and $S \circ \varphi = \varphi^{-1} \circ S$.

Proof. The continuity of φ at points different from a^- and b^- is obvious. Continuity at the two exceptional points requires a little care; if $\sigma = (\text{sign}(\sigma), \sigma_0, \sigma_1, \sigma_2, \dots) \in \Sigma$ agrees with b^- on more than the first 2 coordinates the j_σ of (4.9) exceeds 2. Note that $\sigma_0 = b_0^- = e_j$ and $\text{sign}(\sigma) = \text{sign}(b^-) = -$ by Lemma 4.7. Therefore (4.2) implies that

$$\lim_{t \uparrow 1} (h^{j_\sigma}|_{\sigma_{j_\sigma}})^{-1} \circ \psi_-^{e_j}(t) = \lim_{s \downarrow 0} \left(h^{j_\sigma}|_{\sigma'_{j_\sigma}} \right)^{-1} \circ \psi_-^{e_1}(s),$$

when $\sigma'_{j_\sigma} \in \mathcal{I}^{j_\sigma}$ is determined by (4.10). We conclude from this that $\text{sign}(\sigma') = - = \text{sign}(a^+)$, cf. (4.11). Furthermore, $\sigma'_i = h^{j_\sigma - i}(\sigma'_{j_\sigma}) = a_i^+$ when $i \leq j_\sigma - 2$. Thus we see that $\sigma'_i = a_i^+$ for all $i \leq N$ when $\text{sign}(\sigma) = \text{sign}(b^-)$ and $\sigma_i = b_i^-$ for all $i \leq N + 2$. This proves the continuity of φ at b^- . The continuity at a^- follows in a similar way. To prove that φ is a bijection, we construct the inverse: Let $\sigma = (\text{sign}(\sigma), \sigma_0, \sigma_1, \sigma_2, \dots) \in \Sigma \setminus \{a^+, b^+\}$. We can then consider

$$k_\sigma = \min \left\{ k \in \mathbb{N} : \lim_{s \downarrow 0} (h^k|_{\sigma_k})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right) \neq v \right\}.$$

There is a unique element $\sigma'_{k_\sigma} \in \mathcal{I}^{k_\sigma}$ such that

$$\sigma_{k_\sigma} \cap \sigma'_{k_\sigma} = \lim_{s \downarrow 0} (h^{k_\sigma}|_{\sigma_{k_\sigma}})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right).$$

Set $\sigma' = (\text{sign}(\sigma'), \sigma'_0, \sigma'_1, \sigma'_2, \dots)$, where $\sigma'_i, i < k_\sigma$, is determined by the condition that $\sigma'_i = h^{k_\sigma - i}(\sigma'_{k_\sigma})$ and $\sigma'_i, i \geq k_\sigma$, by the condition that

$$\sigma_i \cap \sigma'_i = \lim_{s \downarrow 0} (h^i|_{\sigma_i})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right).$$

Finally, $\text{sign}(\sigma')$ is determined by the condition that

$$\lim_{t \uparrow 1} \left(h^{k_\sigma}|_{\sigma'_{k_\sigma}} \right)^{-1} \circ \psi_{\text{sign}(\sigma')}^{\sigma'_0}(t) = \lim_{s \downarrow 0} \left(h^{k_\sigma}|_{\sigma_{k_\sigma}} \right)^{-1} \circ \psi_{\text{sign}(\sigma)}^{\sigma_0}(s).$$

We can then define a map $\phi : \Sigma \rightarrow \Sigma$ such that

$$\phi(\sigma) = \begin{cases} \sigma', & \text{when } \sigma \notin \{a^+, b^+\} \\ b^-, & \text{when } \sigma = a^+, \\ a^-, & \text{when } \sigma = b^+. \end{cases}$$

It is straightforward to check that $\varphi \circ \phi = \phi \circ \varphi = \text{id}_\Sigma$, which is all it takes to conclude that φ is a homeomorphism with inverse ϕ . To show that $\varphi \circ S = S \circ \phi$ it suffices to observe that

$$\begin{aligned} \lim_{t \uparrow 1} (h^i|_{\sigma_i})^{-1} \left(\psi_{-\text{sign}(\sigma)}^{\sigma_0}(t) \right) &= \lim_{s \downarrow 0} (h^i|_{\sigma_i})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right) \\ &= \lim_{t \uparrow 1} (h^i|_{\phi(\sigma)_i})^{-1} \left(\psi_{\text{sign}(\phi(\sigma))}^{\phi(\sigma)_0}(t) \right) = \lim_{s \downarrow 0} (h^i|_{\phi(\sigma)_i})^{-1} \left(\psi_{-\text{sign}(\phi(\sigma))}^{\phi(\sigma)_0}(s) \right). \end{aligned}$$

□

Let D_∞ denote the infinite dihedral group, considered here as the semidirect product of $\mathbb{Z}$ and $\mathbb{Z}_2$. Since $\varphi S = S\varphi^{-1}$ it follows that S and φ induce an action α of D_∞ by homeomorphisms of Σ .

Lemma 4.9. α is a free action, i.e. $\alpha_g(\sigma) = \sigma \Rightarrow g = e$.

Proof. Assume that $\sigma \in \Sigma$, $j, k \in \mathbb{Z}$ and that

$$S^j \varphi^k(\sigma) = \sigma. \quad (4.12)$$

Choose $i_0 \in \mathbb{N}$ such that $\#\{I \in \mathcal{I} : I \subseteq e\} > |k|$ for all $i \geq i_0$ and all $e \in \mathbb{E}$. By construction of φ , the intervals $\varphi^l(\sigma)_{i_0}$, $-|k| \leq l \leq |k|$, are mutually distinct. Since $(S^j \varphi^k(\sigma))_{i_0} = \varphi^k(\sigma)_{i_0}$, it follows then from (4.12) that $k = 0$. Once this is established it is clear that $j = 0 \pmod{2}$. □

When $(\sigma, t) \in \Sigma \times [0, 1]$, we define $\Phi_0(\sigma, t) \in \overline{\Gamma}_c$ such that

$$\Phi_0(\sigma, t)_j = \begin{cases} \psi_{\text{sign}(\sigma)}^{\sigma_0}(t), & j = 0, \\ (h^j|_{\sigma_j})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(t) \right), & j \geq 1, 0 < t < 1, \\ \lim_{s \uparrow 1} (h^j|_{\sigma_j})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right), & j \geq 1, t = 1, \\ \lim_{s \downarrow 0} (h^j|_{\sigma_j})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right), & j \geq 1, t = 0. \end{cases}$$

It is straightforward to check that Φ_0 is a continuous surjection. Recall that the *suspension* $\text{susp}(\varphi)$ of φ is the quotient of $\Sigma \times [0, 1]$ obtained by identifying $(\sigma, 1)$ with $(\varphi(\sigma), 0)$. Given a pair $(\sigma, t) \in \Sigma \times [0, 1]$ we denote its image in $\text{susp}(\varphi)$ by $[\sigma, t]$. Since

$\Phi_0(\sigma, 1) = \Phi_0(\varphi(\sigma), 0)$ it follows that Φ_0 falls to a continuous surjection $\Phi : \text{susp}(\varphi) \rightarrow \overline{\Gamma}_c$ defined such that $\Phi[\sigma, t] = \Phi_0(\sigma, t)$.

Since $S\varphi = \varphi^{-1}S$ we can define a homeomorphism $S : \text{susp}(\varphi) \rightarrow \text{susp}(\varphi)$ such that

$$S[\sigma, t] = [S\sigma, 1 - t].$$

S is an involution and we can consider S as an action by $\mathbb{Z}_2$ on $\text{susp}(\varphi)$.

Lemma 4.10. Φ induces a homeomorphism $\Phi : \text{susp}(\varphi)/\mathbb{Z}_2 \rightarrow \overline{\Gamma}_c$.

Proof. It suffices to show that

$$\Phi(\xi) = \Phi(\xi') \Leftrightarrow \xi = \xi' \vee S\xi = \xi'. \quad (4.13)$$

The implication $\Leftarrow$ is obvious. To prove the implication in the other direction, let $\xi = (\sigma, t)$, $\xi' = (\sigma', t')$. If $t \notin \{0, 1\}$ it is clear that $\Phi(\xi) = \Phi(\xi')$ implies that $(\sigma, t) = (\sigma', t')$ or $(\sigma, t) = (S\sigma', 1 - t')$. Assume then that $t = 0$ and that $\Phi(\xi) = \Phi(\xi')$. Then $t' \in \{0, 1\}$. If $\Phi(\xi)_j = v$ for all j , we have that $v \in \sigma_j$ and $\lim_{s \downarrow 0} (h^i|_{\sigma_j})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right) = v$ for all j . It follows then from Lemma 4.7 that $\sigma \in \{a^+, b^+\}$. If also $t' = 0$ we find in the same way that $\sigma' \in \{a^+, b^+\}$. Thus $\xi \in \{[a^+, 0], [b^+, 0]\}$ and $\xi' \in \{[a^+, 0], [b^+, 0]\}$. Since $[a^+, 0] = [\varphi^{-1}(a^+), 1] = [b^-, 1] = S[b^+, 0]$ it follows that $\xi = \xi'$ or $S\xi = \xi'$ in this case. If instead $t' = 1$ we find that $\xi' \in \{[a^-, 1], [b^-, 1]\}$. Since $[a^-, 1] = [\varphi(a^-), 0] = [b^+, 0]$ and $[b^-, 1] = [\varphi(b^-), 0] = [a^+, 0]$ we find again that $\xi = \xi'$ or $S\xi = \xi'$. Assume then that $\Phi(\xi)_j \neq v$ for some j . Then $\lim_{s \downarrow 0} (h^i|_{\sigma_i})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right) = \lim_{s \downarrow 0} (h^i|_{\sigma'_i})^{-1} \left(\psi_{\text{sign}(\sigma')}^{\sigma'_0}(s) \right)$ for all $i \geq j$ (if $t' = 0$) or $\lim_{s \downarrow 0} (h^i|_{\sigma_i})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right) = \lim_{s \uparrow 1} (h^i|_{\sigma'_i})^{-1} \left(\psi_{\text{sign}(\sigma')}^{\sigma'_0}(s) \right)$ for all $i \geq j$ (if $t' = 1$). In the first case, where $t' = 0$, we find that $\varphi S(\sigma) = \sigma'$ and hence that $S\xi = S[\sigma, 0] = S[\varphi^{-1}(\sigma), 1] = [S\varphi^{-1}(\sigma), 0] = [\sigma', 0] = \xi'$ and in the second case, where $t' = 1$, we find that $\sigma = \varphi(\sigma')$ and hence $\xi = [\sigma, 0] = [\varphi(\sigma'), 0] = [\sigma', 1] = \xi'$.

All in all we have now established (4.13) when $t \in [0, 1[$. The case $t = 1$ is handled in way analogues to the case $t = 0$. □

Set $Z = \{(x_i)_{i=0}^\infty \in \overline{\Gamma}_c : x_0 = v\}$ and note that Z is a compact subset of $\overline{\Gamma}_c$, as well as of $W_{\overline{\Gamma}_c, \bar{h}^{-1}}$ with respect to the Wagoner topology. Define $\Lambda : \Sigma \rightarrow Z$ such that

$$\Lambda(\sigma) = \Phi[\sigma, 0].$$

It follows from (4.13) that Λ is a continuous surjection such that

$$\Lambda(\sigma) = \Lambda(\sigma') \Leftrightarrow \sigma' \in \{\sigma, \varphi \circ S(\sigma)\}. \quad (4.14)$$

In particular, Λ is a local homeomorphism which is 2 to 1.

Lemma 4.11. Let $j, k \in \mathbb{Z}$ and let $\sigma \in \Sigma$. There is an open neighborhood U of $\Lambda(\sigma)$ and a local conjugacy (U, V, χ) in Z from $\Lambda(\sigma)$ to $\Lambda(S^j \varphi^k(\sigma))$ such that $\chi(z) = \Lambda(S^j \varphi^k(\Lambda^{-1}(z)))$ for all $z \in U$.

Proof. Note that for every $i \in \mathbb{N}$ the elements $\Lambda(\varphi(\mu))_i$ and $\Lambda(\mu)_i$ of Γ_c are in neighboring intervals from $\mathcal{I}^i$ while $\Lambda(S(\mu))_i$ and $\Lambda(\mu)_i$ are in the same of these intervals. It follows

therefore from (4.8) that $\lim_{i \rightarrow \infty} \sup_{\mu \in \Sigma} d(\Lambda(\varphi(\mu))_i, \Lambda(\mu)_i) = 0$ and $\lim_{i \rightarrow \infty} \sup_{\mu \in \Sigma} d(\Lambda(S(\mu))_i, \Lambda(\mu)_i) = 0$. Hence

$$\lim_{i \rightarrow \infty} \sup_{\mu \in \Sigma} d(\Lambda(S^j \varphi^k(\mu))_i, \Lambda(\mu)_i) = 0.$$

Since Λ is a local homeomorphism there are open neighborhoods U_0 and V_0 of σ and $S^j \varphi^k(\sigma)$ in Σ such that $\Lambda(U_0)$ and $\Lambda(V_0)$ are open in Z and $\Lambda : U_0 \rightarrow \Lambda(U_0)$, $\Lambda : V_0 \rightarrow \Lambda(V_0)$ are homeomorphisms. Set $U = \Lambda(U_0 \cap (S^j \varphi^k)^{-1}(V_0))$. $\square$

Lemma 4.12. *There is a $\delta > 0$ with the following property: When $x, y \in \overline{\Gamma_c}$ are such that $d(x_k, y_k) < \delta$ for all $k \in \mathbb{N}$, it follows that either*

- a) *there is an $N_1 \in \mathbb{N}$ such that x_k and y_k are contained in the closure of the same interval from $\mathcal{I}^1$ for each $k \geq N_1$, or*
- b) *there is an $N_2 \in \mathbb{N}$ such that $x_k, y_k \in J$ for all $k \geq N_2$.*

Proof. In the proof we assume, as we may, that d is an arclength metric. It follows from (4.3) that there is a $\delta_0 > 0$ such that

$$h(\{x \in J_1^+ : d(x, v) \leq \delta_0\}) \subseteq J_1^+$$

and

$$h(\{x \in J_1^- : d(x, v) \leq \delta_0\}) \subseteq J_1^-.$$

Set

$$J_0 = \{x \in J_1^+ : d(x, v) \leq \delta_0\} \cup \{x \in J_1^- : d(x, v) \leq \delta_0\}.$$

Then, if $a \in \{x \in J_1^+ : d(x, v) \leq \delta_0\}$, $b \in \{x \in J_1^- : d(x, v) \leq \delta_0\}$ and $\{h^k(a), h^k(b)\} \not\subseteq J$ for some $k \in \mathbb{N}$, there must be an $l \in \{1, 2, \dots, k\}$ such that $d(h^l(a), h^l(b)) \geq \delta_0$.

There is a $\delta_1 > 0$ such that $d(z, v) < \delta_1 \Rightarrow h(z) \in J_0$ and there is a $\delta_2 > 0$ such that $d(x, y) < \delta_2 \Rightarrow d(h(x), h(y)) < \frac{\delta_1}{2}$. Let $\delta_3 = \min\{d(x, y) : x, y \in h^{-1}(v), x \neq y\}$. Set $\delta = \min\{\delta_0, \delta_1, \delta_2, \delta_3\}$ and consider elements $x, y \in \overline{\Gamma_c}$ such that $d(x_j, y_j) < \delta$ for all j . Assume that there is no N_1 such that a) holds and also no N_2 such that b) holds. There is then an M_0 such that $\{x_{M_0}, y_{M_0}\} \not\subseteq J$, and there is an $M_1 > M_0 + 2$ such that x_{M_1} and y_{M_1} are not both in the closure of the same element from $\mathcal{I}^1$. There is then a path in Γ_c of length $\leq \delta$ which contains an element w from $h^{-1}(v)$ and connects x_{M_1} to y_{M_1} . In particular, it follows that $d(x_{M_1-1}, v) < \delta_1$, $d(y_{M_1-1}, v) < \delta_1$ and hence that $x_{M_1-2}, y_{M_1-2} \in J_0$. If x_{M_1-2}, y_{M_1-2} are both in J_1^+ or both in J_1^- , let $l \geq M_1$ be the least number such that $d(x_l, v) > \delta_1$ or $d(y_l, v) > \delta_1$ which exists because b) fails. There is then an element z from $h^{M_1-l}(v) \setminus \{v\}$ between x_l and y_l such that h^{l-M_1+2} fails to be injective on any open neighborhood of z , contradicting the nonfolding condition. Hence x_{M_1-2} and y_{M_1-2} can not both be in $\{x \in J_1^+ : d(x, v) \leq \delta_0\}$ or both in $\{x \in J_1^- : d(x, v) \leq \delta_0\}$. Since $\{x_{M_0}, y_{M_0}\} \not\subseteq J$, it follows that $d(x_i, y_i) \geq \delta_0$ for some i between M_0 and $M_1 - 2$. This violates the condition that $d(x_j, y_j) < \delta \leq \delta_0$ for all j . $\square$

Lemma 4.13. *Let $\sigma, \sigma' \in \Sigma$. The following are equivalent:*

- i) $\Lambda(\sigma)$ and $\Lambda(\sigma')$ are locally conjugate in Z .
- ii) $\lim_{k \rightarrow \infty} d(\Lambda(\sigma)_k, \Lambda(\sigma')_k) = 0$.
- iii) σ and σ' are in the same D_∞ -orbit under the action α .

Proof. iii) $\Rightarrow$ i) follows from Lemma 4.11 and i) $\Rightarrow$ ii) is obvious. To prove that ii) $\Rightarrow$ iii) assume that $\lim_{i \rightarrow \infty} d(\Lambda(\sigma)_i, \Lambda(\sigma')_i) = 0$. Lemma 4.12 gives us then two alternatives, valid for any $i \geq N$, where N is either the N_1 from a) of Lemma 4.12 or the N_2 from b) of the same lemma: There is an interval, which is either the closure of an interval from $\mathcal{I}^1$ or the interval $J = \psi_-^{e_1}([0, \frac{1}{4}]) \cup \psi_+^{e_j}([0, \frac{1}{4}])$, such that the position of $\Lambda(\sigma)_i$ and $\Lambda(\sigma')_i$ in that interval is

$$\begin{array}{ccccccccccc} & & \Lambda(\sigma)_i & & & & & & & & \Lambda(\sigma')_i & \\ & & \parallel & & & & & & & & \parallel & \\ \text{---} l_1 \text{---} & c_1 & \text{---} l_2 \text{---} & c_2 & \text{---} l_3 \text{---} & c_3 & \text{---} l_4 \text{---} & \dots & \dots & \text{---} l_{k-1} \text{---} & c_{k-1} & \text{---} l_k \text{---} & c_k \end{array}$$

Here the linepieces, which have been labeled l_1 to l_k , are intervals from $\mathcal{I}^i$ and the points labeled c_1 to c_k , with $\Lambda(\sigma)_i$ equal to the first and $\Lambda(\sigma')_i$ equal to the last, are the points from $h^{-i}(v)$ which divide them.

Note that $l_1 = \sigma_i$ or $l_2 = \sigma_i$. Assume first that $\sigma_i = l_1$. Then

$$\Lambda(\sigma)_i = \lim_{s \downarrow 0} (h^i|_{\sigma_i})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right) = \lim_{t \uparrow 1} (h^i|_{\mu_i})^{-1} \left(\psi_{\text{sign}(\mu)}^{\mu_0}(t) \right)$$

where $\mu = \varphi^{-1}(\sigma)$. Hence

$$\Lambda(\varphi^{-1}(\sigma))_i = \lim_{s \downarrow 0} (h^i|_{\mu_i})^{-1} \left(\psi_{\text{sign}(\mu)}^{\mu_0}(s) \right) = c_2.$$

Similarly, $\Lambda(\varphi^{-2}(\sigma))_i = \Lambda(\varphi^{-1}(\varphi^{-1}(\sigma)))_i = c_3$ etc. In particular, $\Lambda(\sigma')_i = \Lambda(\varphi^{-k+1}(\sigma))_i$. Since $i \geq N$, h is injective (and expansive) from an interval containing $\Lambda(\sigma)_{i+1}$ and $\Lambda(\sigma')_{i+1}$ onto an interval containing $l_1, l_2, \dots, l_k$, regardless of which of the two cases from Lemma 4.12 we consider. It follows therefore that $\Lambda(\varphi^{-k+1}(\sigma))_{i+1} = \Lambda(\sigma')_{i+1}$, and by a repetition of the argument $\Lambda(\varphi^{-k+1}(\sigma))_{i'} = \Lambda(\sigma')_{i'}$ for all $i' \geq i$. Since $\Lambda(\varphi^{-k+1}(\sigma))_{i'} = h^{i-i'}(\Lambda(\varphi^{-k+1}(\sigma))_i) = h^{i-i'}(\Lambda(\sigma')_i) = \Lambda(\sigma')_{i'}$ when $i' \leq i$ we conclude that $\Lambda(\varphi^{-k+1}(\sigma)) = \Lambda(\sigma')$. It follows then from (4.14) that σ and σ' are in the same D_∞ -orbit in this case.

Assume next that $\sigma_i = l_2$. Set $\mu = S\varphi^{-1}(\sigma)$ and note that $\Lambda(\mu)_i = l_1$. Furthermore, with $\mu' = \varphi^{-1}(\sigma)$ we find that

$$\begin{aligned} \Lambda(S\varphi^{-1}(\sigma))_i &= \lim_{s \downarrow 0} (h^i|_{\mu_i})^{-1} \left(\psi_{\text{sign}(\mu)}^{\mu_0}(s) \right) = \lim_{t \uparrow 1} (h^i|_{\mu'_i})^{-1} \left(\psi_{\text{sign}(\mu')}^{\mu'_0}(t) \right) \\ &= \lim_{s \downarrow 0} (h^i|_{\sigma_i})^{-1} \left(\psi_{\text{sign}(\sigma)}^{\sigma_0}(s) \right) = \Lambda(\sigma)_i. \end{aligned}$$

We can then argue as above, with σ replaced by $S\varphi^{-1}(\sigma)$, to deduce that $\Lambda(S\varphi^{k-2}(\sigma)) = \Lambda(\varphi^{-k+1}S\varphi^{-1}(\sigma)) = \Lambda(\sigma')$. As above (4.14) tells us then that σ and σ' are in the same D_∞ -orbit. $\square$

Lemma 4.14. *Let (U, V, χ) be a local conjugacy from x to y in Z . Choose $\sigma, \sigma' \in \Sigma$ such that $\Lambda(\sigma) = x$ and $\Lambda(\sigma') = y$. It follows that there is an open and closed neighborhood T of σ and an element $g \in D_\infty$ such that*

- i) $\Lambda(T)$ is open and closed in Z ,
- ii) $\Lambda(T) \subseteq U$,
- iii) $\Lambda : T \rightarrow \Lambda(T)$ is a homeomorphism

- iv) $\alpha_g(\sigma) = \sigma'$, and
- v) $\chi(z) = \Lambda(\alpha_g((\Lambda|_T)^{-1}(z)))$, $z \in \Lambda(T)$.

Proof. Combine i) $\Rightarrow$ iii) of Lemma 4.13 with Lemma 4.11 and Lemma 1.4 of [Th1]. $\square$

Theorem 4.15. *The heteroclinic algebra $B_{\bar{h}^{-1}}(\bar{\Gamma}_c)$ of $(\bar{\Gamma}_c, \bar{h}^{-1})$ is $*$ -isomorphic to $\mathbb{K} \otimes (C(\Sigma) \rtimes_{\alpha} D_{\infty})$.*

Proof. Since α is a free action the crossed product $C(\Sigma) \rtimes_{\alpha} D_{\infty}$ is $*$ -isomorphic to the reduced groupoid C^* -algebra $C_r^*(R)$ where R is the étale equivalence relation given by orbit equivalence under α , with the topology obtained from the product topology of $\Sigma \times D_{\infty}$ via the bijection $(x, g) \rightarrow (x, \alpha_g(x)) \in R$. This follows from [Re] and/or [Ph1]. Since $\Lambda : \Sigma \rightarrow Z$ is a local homeomorphism and Σ is totally disconnected there is an open and closed subset $U \subseteq \Sigma$ such that $\Lambda(U) \subseteq Z$ is open and $\Lambda : U \rightarrow \Lambda(U)$ is a homeomorphism. It follows from Lemma 4.11 and Lemma 4.14 that $\Lambda : U \rightarrow \Lambda(U)$ induces an isomorphism of étale equivalence relations $R|_U \rightarrow R_{\bar{h}^{-1}}(\bar{\Gamma}_c, \Lambda(U))$, where the latter is the étale equivalence relation, for the case at hand, from Section 1.2 of [Th1]. Hence $C_r^*(R|_U)$ is $*$ -isomorphic to $A_{\bar{h}^{-1}}(\bar{\Gamma}_c, \Lambda(U))$, in the notation of [Th1]. Since $B_{\bar{h}^{-1}}(\bar{\Gamma}_c) = A_{\bar{h}^{-1}}(\bar{\Gamma}_c, W_{\bar{\Gamma}_c, \bar{h}^{-1}})$ is simple by Lemma 4.6 it follows from Proposition 4.7 of [Re] that every element of $W_{\bar{\Gamma}_c, \bar{h}^{-1}}$, and in particular every element of Z , is locally conjugate to an element of $\Lambda(U)$. It follows then from Lemma 4.13 that the D_{∞} -orbit of every element σ of Σ contains an element from U . This implies that U is an abstract transversal in R , in the sense of [MRW], so that $C_r^*(R)$ is stably isomorphic to $C_r^*(R|_U)$ by the main result of [MRW]. It follows that $B_{\bar{h}^{-1}}(\bar{\Gamma}_c)$ is stably isomorphic to $C(\Sigma) \rtimes_{\alpha} D_{\infty}$, which gives the theorem since $B_{\bar{h}^{-1}}(\bar{\Gamma}_c)$ is simple by Lemma 4.6. $\square$

Corollary 4.16. *The action α of D_{∞} on Σ is minimal.*

Proof. This was essentially established in the proof of Theorem 4.15, but it follows also from the statement of the theorem by using the wellknown characterisation of ideals in crossed product C^* -algebras of free actions, combined with the simplicity of $B_{\bar{h}^{-1}}(\bar{\Gamma}_c)$, cf. Lemma 4.6. $\square$

Lemma 4.17. *φ is minimal if and only if (Γ_c, h) is not oriented.*

Proof. Let $\psi_e : [0, 1] \rightarrow e$, $e \in \mathbb{E}$, be homeomorphisms determining an orientation of Γ_c . For each pair $e, e' \in \mathbb{E}$, $e \cap h^{-1}(e')$ is the union of closed intervals, $I_1^e, I_2^e, \dots, I_{n(e, e')}^e$, determined by the condition that $h : I_j^e \rightarrow e'$ is a homeomorphism for each j . We will refer to these functions as the characteristic functions of h since they are basically the same as the characteristic functions of Section 3. As observed in the proof of n5) of Section 3 there is a $k \in \mathbb{N}$ and a choice of orientation of Γ_c such that either all the characteristic functions of h^k are increasing or else there are both decreasing and increasing functions among the functions $h^k : I_j^e \rightarrow e'$, $j = 1, 2, \dots, n(e, e')$, for every $e, e' \in \mathbb{E}$. If the latter occurs h can clearly not be oriented. Conversely, if h is not oriented, the first possibility is ruled out by Theorem 3.18 and Lemma 3.6. On the other hand the last condition will hold for all orientations of Γ_c if and only if it holds for one. It follows that what we must show is that φ is minimal if and only if the last condition holds. In

fact, thanks to the mixing condition it suffices to find a $k \in \mathbb{N}$ and a single pair of edges e, e' such that there are both decreasing and increasing functions among the functions $h^k : I_j^e \rightarrow e', j = 1, 2, \dots, n(e, e')$.

Assume first that φ is minimal. Consider the element $a^+ = (-, \text{int } e_1, J_1^-, J_2^-, \dots) \in \Sigma$. Since φ is minimal there is a $M \in \mathbb{N}$ such that $\text{sign}(\varphi^M(a^+)) = +$ and $\varphi^M(a^+)_0 = a_0^+ = \text{int } e_1$. Now take $k_0 \in \mathbb{N}$ so large that e_1 contains more than M intervals from $\mathcal{I}^{k_0}$. It follows then from the definition of φ that the set of functions $h^{k_0} : I_j^{e_1} \rightarrow e_1, j = 1, 2, \dots, n(e_1, e_1)$, can't all have the same orientation; if they did we would have that $\text{sign}(\varphi^M(a^+)) = \text{sign}(a^+) = -$. As pointed out above this gives us the desired conclusion.

(A less direct argument goes as follows: If (Γ_c, h) is oriented, (Γ_c, h^2) is orientable in the sense of [Y4] and it follows then from Corollary 2.7 of [Y4] that $K_1(B_{\bar{h}^{-1}}(\overline{\Gamma_c})) \simeq \mathbb{Z}$. This contradicts Theorem 4.15 and Corollary 4.40. Hence (Γ_c, h) is not oriented.)

Conversely, assume that (Γ_c, h) is not oriented. Thanks to Corollary 4.16 it will be enough to consider an element $\sigma \in \Sigma$ and show that

$$S\sigma \in \overline{\{\varphi^i(\sigma) : i \in \mathbb{Z}\}}.$$

To this end we must consider an $N \in \mathbb{N}$ and show that there is a $j \in \mathbb{Z}$ such that

$$\text{sign}(\varphi^j(\sigma)) = \text{sign}(S\sigma) = -\text{sign}(\sigma) \quad (4.15)$$

and

$$\varphi^j(\sigma)_i = (S\sigma)_i = \sigma_i, \quad i = 0, 1, 2, \dots, N. \quad (4.16)$$

As argued above there is a $k \in \mathbb{N}$ such that there are both decreasing and increasing functions among the functions $h^k : I_j^e \rightarrow e', j = 1, 2, \dots, n(e, e')$, for every $e, e' \in \mathbb{E}$. Let $e, e' \in \mathbb{E}$ be the edges such that $\sigma_N \subseteq e$ and $\sigma_{N+k} \subseteq e'$. By assumption there is a $\sigma' \subseteq e'$ such that $h^k(\sigma') = \sigma_N$ and such that $h^k : \sigma' \rightarrow \sigma_N$ has the opposite orientation of $h^k : \sigma_{N+k} \rightarrow \sigma_N$. Let $l \in \mathbb{N}$ be the number of intervals from $\mathcal{I}^{N+k}$ which sit between σ_{N+k} and σ' in e' . Then $j = l$ or $j = -l$ will satisfy (4.15) and (4.16). $\square$

Theorem 4.15 and Lemma 4.17 will be our main tool for the study of the structure of $B_{\bar{h}^{-1}}(\overline{\Gamma_c})$, but first we pause to show, in the following subsection, how the above double cover and the action of D_∞ is related to the double cover of Fokkink and Yi, cf. [F] and [Y3].

4.3. The relation to the oriented double cover of Yi. This section is not needed for the remaining part of the paper. It is included to put the dihedral action constructed above in perspective, and to prove Proposition 4.24 which may have some independent interest.

Starting from an arbitrary pre-solenoid (not just one which is a wedge of circles) and building on work of R. Fokkink, [F], I. Yi gave in [Y3] an explicit description of an oriented graph Γ' and an orientation preserving continuous map $h' : \Gamma' \rightarrow \Gamma'$ which is a double cover of $h : \Gamma \rightarrow \Gamma$ in the sense that there is 2 to 1 continuous surjection $p : \Gamma' \rightarrow \Gamma$

such that

$$\begin{array}{ccc} \Gamma' & \xrightarrow{h'} & \Gamma' \\ p \downarrow & & \downarrow p \\ \Gamma & \xrightarrow{h} & \Gamma \end{array} \quad (4.17)$$

commutes. See [Y3]. p has the following additional properties:

- 1) $\mathbb{V}_{\Gamma'} = p^{-1}(\mathbb{V}_{\Gamma})$ where $\mathbb{V}_{\Gamma}$ and $\mathbb{V}_{\Gamma'}$ are the vertex sets of Γ and Γ' , respectively.
- 2) For every edge $e \subseteq \Gamma$ there are two edges, e^+ and e^- , such that $p^{-1}(e) = e^+ \cup e^-$ and
 - $p : e^+ \rightarrow e$ is an orientation preserving homeomorphism,
 - $p : e^- \rightarrow e$ is an orientation reversing homeomorphism.

We need a few observations regarding Y 's construction.

Lemma 4.18. *Let $y \in \Gamma$, $z_0 \in \Gamma'$ such that $p(z_0) = h(y)$. Then*

$$\#\{z \in p^{-1}(y) : h'(z) = z_0\} = 1.$$

Proof. Assume first that $x = p(z_0) \notin \mathbb{V}_{\Gamma}$. There are then edges $e, f \subseteq \Gamma$ and an open non-degenerate interval $I \subseteq e$ such that $y \in I$ and $h(I) = \text{int } f \ni x$. It follows from 1) and 2) that $p^{-1}(y) = \{z_+, z_-\}$ where $z_{\pm} \in \text{int } e_{\pm}$. Similarly, $z_0 \in \text{int } f_+$ or $z_0 \in \text{int } f_-$. Assume that $z_0 \in \text{int } f_+$. If $h : I \rightarrow f$ is orientation preserving the commutative of (4.17) implies that $\{z \in p^{-1}(y) : h'(z) = z_0\} = \{z_+\}$ since h' is orientation preserving. If instead $h : I \rightarrow f$ is orientation reversing we conclude in the same way that $\{z \in p^{-1}(y) : h'(z) = z_0\} = \{z_-\}$. Similarly, when $z_0 \in \text{int } f_-$ we find that $\{z \in p^{-1}(y) : h'(z) = z_0\} = \{z_+\}$ when $h : I \rightarrow f$ is orientation reversing and $\{z \in p^{-1}(y) : h'(z) = z_0\} = \{z_-\}$ when it is orientation preserving.

To handle the case where $x \in \mathbb{V}_{\Gamma}$, let $p^{-1}(y) = \{z_1, z_2\}$. Assume to reach a contradiction that $h'(z_1) = h'(z_2) = z_0$. Use then that p is 2 to 1 and hence a local homeomorphism to find open neighborhoods U and V of y and x , respectively, together with open sets $Z_i \subseteq \Gamma'$, $W_i \subseteq \Gamma'$, $i = 1, 2$, such that $Z_1 \cap Z_2 = \emptyset$, $z_0 \in W_1$, $z_i \in Z_i$, $i = 1, 2$, $W_1 \cap W_2 = \emptyset$, $p(Z_1) = p(Z_2) = U$, $h(U) \subseteq V$ and $p(W_1) = p(W_2) = V$, and such that p is injective on W_1 . Since $h'(z_1) = h'(z_2) = z_0$ we can shrink U (and hence Z_i , $i = 1, 2$) to arrange that $h'(Z_i) \subseteq W_1$, $i = 1, 2$. For every element ξ of U there is an element $a_i \in Z_i$ such that $p(a_i) = \xi$, $i = 1, 2$. Note that $h'(a_i) \in W_1$ and that $ph'(a_i) = hp(a_i) = h(\xi)$, $i = 1, 2$. Since p is injective on W_1 it follows that $h'(a_1) = h'(a_2)$. This contradicts the first part of the proof provided $h(\xi) \notin \mathbb{V}_{\Gamma}$. Since h does not collapse U to a finite set there is a $\xi \in U$ such that $h(\xi) \notin \mathbb{V}_{\Gamma}$. This shows that $\#\{z \in p^{-1}(y) : h'(z) = z_0\} \leq 1$. Finally, the conclusion that $\#\{z \in p^{-1}(y) : h'(z) = z_0\} > 0$ follows by approximating y with elements y' for which $h(y') \notin \mathbb{V}_{\Gamma}$. $\square$

Define $\bar{p} : \bar{\Gamma}' \rightarrow \bar{\Gamma}$ such that $\bar{p}((x_i)_{i=0}^{\infty}) = (p(x_i))_{i=0}^{\infty}$. It follows from (4.17) that $\bar{p} \circ \bar{h} = \bar{h}' \circ \bar{p}$.

Lemma 4.19. *Let $x, y \in \bar{\Gamma}'$ such that $\bar{p}(x) = \bar{p}(y)$. Then $x = y$ if and only if $x_0 = y_0$.*

Proof. This follows immediately from Lemma 4.18. $\square$

Lemma 4.20. *Let $\{x^n\}_{n=1}^\infty$ be a sequence in $\overline{\Gamma'}$ and $x \in \overline{\Gamma'}$ an element in $\overline{\Gamma'}$. Then $\lim_{n \rightarrow \infty} x^n = x$ if and only if $\lim_{n \rightarrow \infty} \bar{p}(x^n) = \bar{p}(x)$ and $\lim_{n \rightarrow \infty} x_0^n = x_0$.*

Proof. One implication follows from the continuity of $\bar{p}$. To prove the other assume that $\lim_{n \rightarrow \infty} \bar{p}(x^n) = \bar{p}(x)$ and $\lim_{n \rightarrow \infty} x_0^n = x_0$. If x^n does not converge to x there is a subsequence $\{x^{n_i}\}$ and an element $y \neq x$ such that $\lim_{i \rightarrow \infty} x^{n_i} = y$. But then the assumptions imply that $\bar{p}(y) = \bar{p}(x)$ and $x_0 = y_0$. This contradicts Lemma 4.19. $\square$

To relate Yi's double cover to the constructions of the preceding section we have to specialize, as we did above, to the case where Γ_c is a wedge of circles. Let $(\sigma, t) \in \Sigma \times [0, 1]$. It follows from Lemma 4.18 and Lemma 4.19 that there is unique element $\Psi_0(\sigma, t) \in \overline{\Gamma'_c}$ such that $\bar{p}(\Psi_0(\sigma, t)) = \Phi_0(\sigma, t)$ and

$$\Psi_0(\sigma, t)_0 \in \sigma_0^{\text{sign}(\sigma)}.$$

Since $\Phi_0(\varphi(\sigma), 0) = \Phi_0(\sigma, 1)$, it follows from Lemma 4.19 that $\Psi_0(\sigma, 1) = \Psi_0(\varphi(\sigma), 0)$. We get therefore a well-defined map

$$\Psi : \text{susp}(\varphi) \rightarrow \overline{\Gamma'_c}$$

defined such that $\Psi[\sigma, t] = \Psi_0(\sigma, t)$. Then the diagram

$$\begin{array}{ccc} \text{susp}(\varphi) & \xrightarrow{\Psi} & \overline{\Gamma'_c} \\ & \searrow \Phi & \downarrow \bar{p} \\ & & \overline{\Gamma_c} \end{array} \quad (4.18)$$

commutes by construction.

Lemma 4.21. $\Psi : \text{susp}(\varphi) \rightarrow \overline{\Gamma'_c}$ is a homeomorphism.

Proof. It follows from Lemma 4.20 that Ψ_0 is continuous and the same is therefore true for Ψ . Let $x \in \overline{\Gamma'_c}$. Since Φ is surjective there is an element $(\sigma, t) \in \Sigma \times [0, 1]$ such that $\bar{p}(x) = \Phi(\sigma, t)$. Then $x_0 \in \sigma_0^{\text{sign}(\sigma)}$ or $x_0 \in \sigma_0^{-\text{sign}(\sigma)}$. In the first case, $x = \Psi_0(\sigma, t)$ and in the second $\Psi_0(S\sigma, 1 - t) = x$. This shows that Ψ is surjective. Thanks to this it follows from (4.18) and the fact that both $\bar{p}$ and Φ are 2 to 1 that Ψ is also injective. $\square$

To see how the D_∞ -action fits into this picture, note that we can also consider $\text{susp}(\varphi)$ as the quotient of $\Sigma \times \mathbb{R}$ under the equivalence relation $\sim$ where $(\sigma, t) \sim (\sigma', t')$ means that $t - t' \in \mathbb{Z}$ and $\varphi^{t-t'}(\sigma) = \sigma'$. We obtain then a flow $\varphi_t, t \in \mathbb{R}$, on $\text{susp}(\varphi)$ defined such that

$$\varphi_s[\sigma, t] = [\sigma, t + s].$$

Note that $S \circ \varphi_t = \varphi_{-t} \circ S$ so that we have in fact an action β of the generalized dihedral group $D_\mathbb{R} = \mathbb{R} \rtimes \mathbb{Z}_2$ on $\text{susp}(\varphi)$. Formally $D_\mathbb{R}$ is the semidirect product $\mathbb{R} \rtimes \mathbb{Z}_2$ in the category of locally compact groups coming from the automorphism $t \mapsto -t$ of $\mathbb{R}$. If we write the elements of $D_\mathbb{R}$ as tuples (i, s) where $s \in \mathbb{R}$ and $i \in \mathbb{Z}_2 = \{0, 1\}$,

$$\beta_{(i,s)}[\sigma, t] = S^i \varphi_s[\sigma, t] = \left[(\varphi S)^i \sigma, (-1)^i (s + t) \right].$$

Lemma 4.22. β is a free action, i.e. $\beta_x(\xi) = \xi \Rightarrow x = 0$.

Proof. If $x = (i, s)$ and $\beta_{(i,s)}[\sigma, t] = [(\varphi S)^i \sigma, (-1)^i(s+t)] = [\sigma, t]$, where $t \in [0, 1]$, we have that

$$\left[\varphi^{[(-1)^i(s+t)]+i} S^i \sigma, r \right] = [\sigma, t], \quad (4.19)$$

where $[(-1)^i(s+t)]$ denotes the integer part of $(-1)^i(s+t)$ and $r = (-1)^i(s+t) - [(-1)^i(s+t)]$. It follows from (4.19) that $r = t$ and $\varphi^{[(-1)^i(s+t)]+i} S^i \sigma = \sigma$ or $r = 0$ and $t = 1$. In the first case we deduce from Lemma 4.9

$$[(-1)^i(s+t)] + i = 0, \quad (-1)^i = 1, \quad r = t.$$

It follows that $x = 0$ in this case. In the second case we conclude from Lemma 4.9 that

$$[(-1)^i(s+t)] + i - 1 = 0, \quad (-1)^i = 1$$

which shows that $x = 0$, also in this case. $\square$

Lemma 4.23. *Let $x, y \in \overline{\Gamma}'_c$. Then $\lim_{n \rightarrow \infty} d(\overline{h}^{-n}(\Phi(x)), \overline{h}^{-n}(\Phi(y))) = 0$ if and only if $\beta_a(x) = y$ for some $a \in D_{\mathbb{R}}$.*

Proof. Assume first that $\beta_a(x) = y$ for some $a \in D_{\mathbb{R}}$. Write $a = (i, t)$ where $t \in \mathbb{R}$ and $i \in \mathbb{Z}_2$. Since $\Phi \circ S = \Phi$ we may assume that $i = 0$ so that $\varphi_t(x) = y$. Write $x = (\sigma, s)$ where $s \in [0, 1]$. Then $\varphi_t(x) = [\varphi^k(\sigma), r]$ where $k \in \mathbb{Z}$ and $r \in [0, 1]$. Since $\Phi([\varphi^k(\sigma), r])_n$ and $\Phi([\varphi^k(\sigma), 0])_n$ are in the same interval from $\mathcal{I}^n$ for each n it follows from (4.8) that $\lim_{n \rightarrow \infty} d(\overline{h}^{-n}(\Phi([\varphi^k(\sigma), r])), \overline{h}^{-n}(\Phi([\varphi^k(\sigma), 0]))) = 0$. By Lemma 4.11 $\lim_{n \rightarrow \infty} d(\overline{h}^{-n}(\Phi([\varphi^k(\sigma), 0])), \overline{h}^{-n}(\Phi([\sigma, 0]))) = 0$, and by a repetition of the preceding argument $\lim_{n \rightarrow \infty} d(\overline{h}^{-n}(\Phi([\sigma, 0])), \overline{h}^{-n}(\Phi([\sigma, s]))) = 0$, proving that $\lim_{n \rightarrow \infty} d(\overline{h}^{-n}(\Phi(\varphi_t(x))), \overline{h}^{-n}(\Phi(x))) = 0$, as dsired.

Assume next that $\lim_{n \rightarrow \infty} d(\overline{h}^{-n}(\Phi(x)), \overline{h}^{-n}(\Phi(y))) = 0$. Write $x = [\sigma, t]$ and $y = [\sigma', t']$ where $\sigma, \sigma' \in \Sigma, t, t' \in [0, 1]$. Since $[\sigma, t] = \varphi_t[\sigma, 0]$ and $[\sigma', t'] = \varphi_{t'}[\sigma', 0]$ and $\lim_{n \rightarrow \infty} d(\overline{h}^{-n}(\Phi(\varphi_{-t}(x))), \overline{h}^{-n}(\Phi(\varphi_{-t'}(y)))) = 0$ by the first part of the proof we may assume that $t = t' = 0$, and then the desired conclusion follows from Lemma 4.13. $\square$

Proposition 4.24. *Let Γ be a pre-solenoid and Γ' the corresponding oriented double cover of Γ . Let $\overline{p}: \overline{\Gamma}' \rightarrow \overline{\Gamma}$ be the corresponding canonical surjection. There is then a free action β of $D_{\mathbb{R}}$ on $\overline{\Gamma}'$ with the following property:*

Two elements $x, y \in \overline{\Gamma}'$ satisfy that

$$\lim_{n \rightarrow \infty} d(\overline{h}^{-n}(\overline{p}(x)), \overline{h}^{-n}(\overline{p}(y))) = 0$$

if and only if $\beta_a(x) = y$ for some $a \in D_{\mathbb{R}}$.

Proof. When Γ is a wedge of circles the existence follows from the preceding. For the general case we use a result of R. Williams which shows that for an arbitrary pre-solenoid (Γ, h) there is an $m \in \mathbb{N}$ and a pre-solenoid (Γ_1, h_1) such that Γ_1 is a wedge of circles and $(\overline{\Gamma}, \overline{h}^m)$ is conjugate to $(\overline{\Gamma}_1, \overline{h}_1)$, cf. Proposition 2.2 of [Y4]. Since this conjugacy lifts to

a homeomorphism between the double covers $\overline{\Gamma'}$ and $\overline{\Gamma'_1}$ by Proposition 3.7 of [Y3] this gives us the general case. $\square$

Remark 4.25. It follows from Corollary 4.16 that the action β of Proposition 4.24 is minimal. It is a consequence of Proposition 4.27 which we prove below that the action is also uniquely ergodic.

4.4. Affability and a unique trace. In this section we obtain the conclusion that $B_{\overline{\Gamma}}^{-1}(\overline{\Gamma})$ has a densely defined lower-semicontinuous trace which is unique up to multiplication by a scalar. For this purpose we return to the setting of Section 4.2. In particular, we consider the intervals $\mathcal{J}$ given by (4.1) and use them to define an ordered Bratteli diagram $\mathbb{B}$ as follows: The first level has $2n$ vertices, labeled $(+, i)$ and $(-, i)$, $i = 1, 2, \dots, n$. Beyond this first level the diagram is stationary and has n vertices at each level and there are a_{ik} arrows, labeled by I_{ik}^l , $l = 1, 2, \dots, a_{ik}$, from vertex k at each level to vertex i at the next level. From the first to the second level there are a_{ik} arrows, labeled by I_{ik+}^l , $l = 1, 2, \dots, a_{ik}$, from vertex $(+, k)$ at the first level to vertex i at the next level, and there are a_{ik} arrows, labeled by I_{ik-}^l , $l = 1, 2, \dots, a_{ik}$, from vertex $(-, k)$ at the first level to vertex i at the next level. Then every element $\sigma \in \Sigma$ defines an infinite path $\xi(\sigma) = (I_{i_0 j_0 \text{sign}(\sigma)}^{l_0}, I_{i_1 j_1}^{l_1}, I_{i_2 j_2}^{l_2}, \dots)$ in $\mathbb{B}$ by the requirement that $\sigma_{k+1} \subseteq I_{i_k j_k}^{l_k}$, $k \geq 0$. Let $X_{\mathbb{B}}$ be the space of infinite paths in $\mathbb{B}$ equipped with the relative topology inherited from the infinite product space $\{I_{ik}^l : l = 1, 2, \dots, a_{ik}, i, k = 1, 2, \dots, n\}^{\mathbb{N}}$. It is straightforward to check that $\xi : \Sigma \rightarrow X_{\mathbb{B}}$ is a homeomorphism.

Consider a vertex w in $\mathbb{B}$, say the vertex i at some level beyond the second level of the Bratteli diagram. Then the arrows in $\mathbb{B}$ terminating at w are labeled by I_{il}^k , $k = 1, 2, \dots, a_{il}^k$, $l = 1, 2, \dots, n$. We equip these arrows with a total ordering such that I_{i1}^1 is the minimal element and $I_{ij}^{a_{ij}}$ the maximal. If $w = i$ is at the second level the arrows in $\mathbb{B}$ terminating at w are labeled by $I_{il\pm}^k$, $k = 1, 2, \dots, a_{il}^k$, $l = 1, 2, \dots, n$. We equip these arrows with a total ordering such that I_{i1-}^1 is the minimal element and $I_{ij+}^{a_{ij}}$ the maximal. Add a zeroth level with one vertex with $2n$ emitting arrows, one to each of the vertices at the first level. With these additional features $\mathbb{B}$ becomes a simple ordered Bratteli diagram in the sense of [GPS1] and by construction there is a unique minimal and a unique maximal path in $X_{\mathbb{B}}$. Hence the Bratteli-Vershik map $\psi : X_{\mathbb{B}} \rightarrow X_{\mathbb{B}}$ is defined and minimal. For $x \in X_{\mathbb{B}}$ and $\sigma \in \Sigma$ we let $\text{orb}_{\psi}(x)$ denote the orbit of x under ψ and $\text{orb}_{D_{\infty}}(\sigma)$ the orbit of σ under α .

Lemma 4.26. $\xi(\text{orb}_{D_{\infty}}(\sigma)) = \text{orb}_{\psi}(\xi(\sigma))$ for all $\sigma \in \Sigma$.

Proof. Assume $\sigma' \in \text{orb}_{D_{\infty}}(\sigma)$. It follows from the implication iii) $\Rightarrow$ ii) of Lemma 4.13 that $\lim_{j \rightarrow \infty} \text{dist}(\sigma_j, \sigma'_j) = 0$. By Lemma 4.12 this implies that there is an $N \in \mathbb{N}$ such that either $\xi(\sigma)_k = \xi(\sigma')_k$ for all $k \geq N$ or else $\xi(\sigma)_k \neq \xi(\sigma')_k$ and $\{\xi(\sigma)_k, \xi(\sigma')_k\} = \{I_{11}^1, I_{jj}^{a_{jj}}\}$ for all $k \geq N$. In both cases we conclude that $\xi(\sigma') \in \text{orb}_{\psi}(\xi(\sigma))$, cf. [GPS1], in particular Remark (ii) on page 72 of [GPS1].

Conversely, if $\xi(\sigma') \in \text{orb}_{\psi}(\xi(\sigma))$, it follows from [GPS1] that there is an $N \in \mathbb{N}$ such that either $\xi(\sigma)_k = \xi(\sigma')_k$ for all $k \geq N$ or else $\xi(\sigma)_k \neq \xi(\sigma')_k$ and $\{\xi(\sigma)_k, \xi(\sigma')_k\} = \{I_{11}^1, I_{jj}^{a_{jj}}\}$ for all $k \geq N$. In both cases we conclude that $\lim_{k \rightarrow \infty} d(\Lambda(\sigma)_k, \Lambda(\sigma')_k) = 0$ and then Lemma 4.13 implies that $\sigma' \in \text{orb}_{D_{\infty}}(\sigma)$. $\square$

It follows from Lemma 4.26 that the étale equivalence relation given by orbit equivalence under α is affable in the sense of [GPS2]. For the purposes of the present paper it is the following consequence of Proposition 4.26 we shall need:

Proposition 4.27. *The action α of D_∞ on Σ has only one invariant Borel probability measure, and hence $C(\Sigma) \rtimes_\alpha D_\infty$ has a unique trace state.*

Proof. The simplex of ψ -invariant Borel probability measures on $X_{\mathbb{B}}$ agrees with the trace-simplex of the unital AF-algebras whose Bratteli-diagram is $\mathbb{B}$, cf. Theorem 1.13 and Theorem 3.7 of [GPS1]. But $\mathbb{B}$ is eventually stationary with an inclusion pattern given by a primitive matrix and hence this AF-algebra has a unique trace state by [Ef]. In this way the first statement becomes a well-known consequence of Lemma 4.26 and the second is then a well-known consequence of the first, cf. e.g. Theorem 4.5 of [Th2]. $\square$

It follows from Proposition 4.27 and Theorem 4.15 that the heteroclinic algebra $B_{h^{-1}}(\overline{\Gamma}) \simeq B_{h^{-1}}(\overline{\Gamma}_c)$ has an essentially unique densely defined lower-semicontinuous trace. The remaining information about the structure which we shall obtain in this paper comes from the fact that it is (stably) given as the crossed product of a free minimal action of D_∞ on the Cantor set. The next two sections are therefore devoted to a study of the crossed products by such actions.

4.5. On crossed products $C(\Omega) \rtimes D_\infty$. An action α of the infinite dihedral group D_∞ on the Cantor set Ω is generated by two homeomorphism φ and σ of Ω such that

$$\sigma^2 = \text{id} \tag{4.20}$$

and

$$\sigma\varphi\sigma = \varphi^{-1}. \tag{4.21}$$

Crossed product C^* -algebras arising from actions of D_∞ on Ω were studied by Bratteli, Evans and Kishimoto in [BEK] under the assumptions that φ is minimal and that σ or $\varphi\sigma$ has a fixed point. The main result of [BEK] is that the crossed product is then AF. As we have seen, actions of D_∞ on Ω arising from 1-solenoids are always free so the last assumption in [BEK] is never met for such actions. Minimality of φ , on the other hand, is equivalent to orientability of the 1-solenoid, as it follows by combining Lemma 4.17 with Theorem 2.2.

We will assume throughout, in this and the following section, that α is a free and minimal action; i.e. for every $x \in \Omega$ the points $\varphi^j(x), \sigma\varphi^k(x), j, k \in \mathbb{Z}$, are all different and constitute a dense subset of Ω .

Lemma 4.28. *One of the following two alternatives occur:*

- i) φ is minimal on Ω or
- ii) there is a clopen φ -invariant subset $F \subseteq \Omega$ such that $\varphi|_F$ is minimal (on F), $\sigma(F) \cap F = \emptyset$ and $\sigma(F) \cup F = \Omega$.

If F' is another subset of Ω with the properties specified in ii), $F' = F$ or $F' = \sigma(F)$.

Proof. Assume that φ is not minimal. There is then a non-trivial closed subset $F \subseteq \Omega$ such that $\varphi(F) = F$. Since $F \cap \sigma(F)$ is closed and invariant under both φ and σ , the minimality of α implies that $F \cap \sigma(F) = \emptyset$ or $F \cap \sigma(F) = \Omega$. The latter possibility can not occur since F is non-trivial, i.e. $F \cap \sigma(F) = \emptyset$. A similar reasoning, with $F \cup \sigma(F)$

replacing $F \cap \sigma(F)$, shows that $F \cup \sigma(F) = \Omega$. In particular, F is clopen. If F_0 is any non-empty closed φ -invariant subset of F the same arguments show that $F_0 \cap \sigma(F_0) = \emptyset$ and $F_0 \cup \sigma(F_0) = \Omega$. It follows that F_0 must be all of F , and hence $\varphi|_F$ is minimal. This proves the existence of F . The essential uniqueness of F , as described in the statement, is established along similar lines. $\square$

Let A and B be isomorphic C^* -algebras, and let $\gamma : A \rightarrow B$ be a $*$ -isomorphism. The automorphism of $A \oplus B$ sending (a, b) to $(\gamma^{-1}(b), \gamma(a))$ will be called a *swop* of A .

Lemma 4.29. *Let $\gamma : A \oplus B \rightarrow A \oplus B$ be a swop of A . Then*

$$(A \oplus B) \rtimes_{\gamma} \mathbb{Z}_2 \simeq M_2(A).$$

Proof. Left to the reader. $\square$

Theorem 4.30. *Assume that φ is not minimal, and let F be the clopen subset from ii) of Lemma 4.28. Then*

$$C(\Omega) \rtimes_{\alpha} D_{\infty} \simeq M_2(C(F) \rtimes_{\varphi} \mathbb{Z}), \quad (4.22)$$

and hence $C(\Omega) \rtimes_{\alpha} D_{\infty}$ is a simple unital AT-algebra of real rank zero with K_1 -group $\mathbb{Z}$.

Proof. Since both F and $\sigma(F)$ are φ -invariant we have a decomposition

$$C(\Omega) \rtimes_{\varphi} \mathbb{Z} \simeq (C(F) \rtimes_{\varphi} \mathbb{Z}) \oplus (C(\sigma(F)) \rtimes_{\varphi} \mathbb{Z}).$$

This isomorphism turns σ into the swop of $C(F) \rtimes_{\varphi} \mathbb{Z}$ given by the $*$ -isomorphism $C(F) \rtimes_{\varphi} \mathbb{Z} \rightarrow C(\sigma(F)) \rtimes_{\varphi} \mathbb{Z}$ which sends $f \in C(F)$ to $f \circ \sigma \in C(\sigma(F))$ and u to u^* when u denotes the canonical unitary going with a crossed product by $\mathbb{Z}$. Hence Lemma 4.29 applies. The last statement follows from (4.22) and [HPS]. $\square$

We turn to the case where φ is minimal.

Proposition 4.31. *Assume that φ is minimal. Let $R \in \mathbb{N}$ be given. There is then a partition*

$$C(k, i), \quad i = 0, 1, 2, \dots, J_k - 1, \quad k = 1, 2, \dots, N,$$

of Ω into clopen sets such that

- i) $J_k \geq R, k = 1, 2, \dots, N,$
- ii) $\sigma(C(k, i)) = C(k, J_k - i - 1)$ for all $i, k,$
- iii) $\varphi(C(k, i)) = C(k, i + 1), i = 0, 1, \dots, J_k - 2,$ for all $k,$
- iv) $\varphi\left(\bigcup_{k=1}^N C(k, J_k - 1)\right) = \bigcup_{k=1}^N C(k, 0).$

Proof. Since D_{∞} acts freely, any point in Ω has a clopen neighborhood Z such that

$$[\varphi^k(Z) \cup \varphi^k(\varphi\sigma(Z))] \cap [\varphi^l(Z) \cup \varphi^l(\varphi\sigma(Z))] = \emptyset$$

when $k \neq l$ and $0 \leq k, l \leq R$. The desired partition can then be obtained by applying Lemma 1.4 of [BEK] with $Y = Z \cup \varphi\sigma(Z)$. $\square$

Theorem 4.32. *Assume that φ is minimal. The action of $\mathbb{Z}_2$ on $C(\Omega) \rtimes_{\varphi} \mathbb{Z}$ given by σ has the tracial Rokhlin property of Phillips [Ph4].*

Proof. By [HPS] $C(\Omega) \rtimes_{\varphi} \mathbb{Z}$ is a unital simple AT-algebra of real rank zero. It follows therefore from (2.21) of [EG] that $C(\Omega) \rtimes_{\varphi} \mathbb{Z}$ has tracial rank zero. By Theorem 2.9 of [Ph5] the conclusion will follow if we for a given $\epsilon > 0$ exhibit functions $f_1, f_2 \in C(\Omega)$ such that $0 \leq f_i \leq 1$, $\|f_i - f_i \circ \varphi\| \leq \epsilon$, $i = 1, 2$, $f_1 f_2 = 0$, $f_1 \circ \sigma = f_2$ and $\omega(1 - f_1 - f_2) \leq \epsilon$ for every tracial state ω of $C(\Omega) \rtimes_{\varphi} \mathbb{Z}$. To this end, choose first an $M \in \mathbb{N}$ such that $\frac{1}{M} \leq \epsilon$ and then an $R \geq M$ such that $\frac{4M}{R} \leq \epsilon$. With this choice of R , consider a partition $C(k, i)$ with the properties spelled out in Proposition 4.31. For each $k = 1, 2, \dots, N$ we define a function $g_k : \Omega \rightarrow [0, 1]$ in the following way: Let l_k be the integer part of $\frac{J_k - 1}{2}$. Choose numbers $\alpha_i \in [0, 1]$, $i = 0, 1, 2, \dots, l_k$, such that $\alpha_0 = \alpha_{l_k} = 0$, $|\alpha_{j+1} - \alpha_j| \leq \epsilon$ for all $j = 0, 1, 2, \dots, l_k - 1$, and $\alpha_i = 1$, $M \leq i \leq l_k - M - 1$. This is possible since $\frac{1}{M} \leq \epsilon$. Now define g_k to be constant equal to α_i on $C(k, i)$ and to be zero on $\Omega \setminus \left(\bigcup_{i=0}^{l_k} C(k, i)\right)$.

It is then easy to check that $\|g_k \circ \varphi - g_k\| \leq \epsilon$ and $\|g_k \circ \varphi^{-1} - g_k\| \leq \epsilon$, and that $g_k \circ \varphi$, $g_k \circ \varphi^{-1}$ are both supported in $\bigcup_{i=0}^{l_k} C(k, i)$. Set

$$f_1 = \sum_{k=1}^N g_k$$

and $f_2 = f_1 \circ \sigma$. It is then easy to check that $0 \leq f_i \leq 1$, $\|f_i - f_i \circ \varphi\| \leq \epsilon$, $i = 1, 2$, $f_1 f_2 = 0$, $f_1 \circ \sigma = f_2$. To check that $\omega(1 - f_1 - f_2) \leq \epsilon$ for every tracial state ω of $C(\Omega) \rtimes_{\varphi} \mathbb{Z}$, observe first that every D_{∞} -invariant trace state is given by integration with respect to a D_{∞} -invariant Borel probability measure on Ω ; this follows e.g. from a more general result which can be found as Theorem 4.5 of [Th2]. Hence it suffices to show that

$$\int_{\Omega} 1 - f_1 - f_2 \, d\mu \leq \epsilon \quad (4.23)$$

when μ is a D_{∞} -invariant Borel probability measure on X . To this end note that

$$\int_{\Omega} 1 - f_1 - f_2 \, d\mu \leq \sum_{k=1}^N 4M\beta_k$$

where $\beta_k = \mu(C(k, 0))$. Since $\frac{4M}{J_k} \leq \frac{4M}{R} \leq \epsilon$ and $\sum_{k=1}^N J_k \beta_k = 1$ this yields (4.23). $\square$

Corollary 4.33. *Assume that φ is minimal. Then $C(\Omega) \rtimes_{\alpha} D_{\infty}$ has tracial rank zero in the sense of Lin, [Lin2].*

Proof. This follows from Theorem 4.32 and Theorem 2.6 of [Ph4]. $\square$

Remark 4.34. The $\mathbb{Z}_2$ action of Theorem 4.32 does not have the (strict) Rokhlin property. Indeed, if it did the crossed product $C(\Omega) \rtimes_{\alpha} D_{\infty}$ would be a simple AT-algebra by Corollary 3.4 of [OP] which is not possible since we shall show below that the K_0 -group of $C(\Omega) \rtimes_{\alpha} D_{\infty}$ contains torsion.

It follows *almost* from the preceding theorem that $C(\Omega) \rtimes_{\alpha} D_{\infty}$ is classified by K-theory, thanks to the work of H. Lin. What is missing is that we need to show that $C(\Omega) \rtimes_{\alpha} D_{\infty}$ satisfies the UCT. This technicality is dealt with in the following proposition.

Definition 4.35. A separable C^* -algebra A satisfies the UCT when the canonical map

$$\gamma_{A,D} : KK(A, D) \rightarrow \text{Hom}(K_*(A), K_*(D))$$

is an isomorphism for every separable C^* -algebra D such that $K_0(D)$ and $K_1(D)$ are both divisible groups.

That this formulation is equivalent to the more conventional, which appears in the literature on the classification of C^* -algebras, follows from Proposition 23.8.1 of [Bl].

Proposition 4.36. *Let A_1 and A_2 be a separable C^* -algebras with a common C^* -subalgebra B . Assume that A_1 and A_2 satisfy the UCT, and assume that B is AF. Then the amalgamated free product $A_1 *_B A_2$ satisfies the UCT.*

Proof. Note that $A_1 *_B A_2 = \varinjlim_n A_1 *_B A_2$ when $B_1 \subseteq B_2 \subseteq B_3 \subseteq \dots$ is an increasing sequence of finite-dimensional C^* -subalgebras with dense union in B . As shown in [Th3] $A_1 *_B A_2$ is KK-equivalent to the mapping cone for an embedding of B_n into $A_1 \oplus A_2$. Therefore $A_1 *_B A_2$ satisfies the UCT since A_1 , A_2 and B_n all do. Since the class of separable C^* -algebras which satisfy the UCT is closed under inductive limits, cf. [Bl], it follows that $A_1 *_B A_2$ is in the class. $\square$

Corollary 4.37. $C(\Omega) \rtimes_\alpha D_\infty$ satisfies the UCT.

Proof. As observed and used in Section 4 of [BEK], $C(\Omega) \rtimes_\alpha D_\infty$ can be realized as the amalgamated free product

$$(C(\Omega) \rtimes_\sigma \mathbb{Z}_2) *_C (C(\Omega) \rtimes_{\varphi\sigma} \mathbb{Z}_2).$$

Both $C(\Omega) \rtimes_\sigma \mathbb{Z}_2$ and $C(\Omega) \rtimes_{\varphi\sigma} \mathbb{Z}_2$ are sub-homogeneous and type I (see Section 4 of [BEK] or (4.25) below), and $C(\Omega)$ is AF, so Proposition 4.36 applies. $\square$

We now turn to the calculation of $K_*(C(\Omega) \rtimes_\alpha D_\infty)$. When φ is not minimal Theorem 4.30 shows that $C(\Omega) \rtimes_\alpha D_\infty$ is, up to an M_2 -tensor factor, also the crossed product coming from a minimal homeomorphism of the Cantor set, and an obvious construction shows that all such algebras arise this way. In particular, the K -theory can be found from [HPS]; the K_1 -group is $\mathbb{Z}$ and the K_0 -group is a simple dimension group. Also from [HPS] we conclude that all simple dimension groups occur here.

It remains to consider the case where φ is minimal. It follows then from [N] and Lemma 4.2 of [BEK] that $K_0(C(\Omega) \rtimes_\varphi \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2)$ is the cokernel of the map

$$K_0(C(\Omega)) \xrightarrow{(i_{1*}, i_{2*})} K_0(C(\Omega) \rtimes_\sigma \mathbb{Z}_2) \oplus K_0(C(\Omega) \rtimes_{\varphi\sigma} \mathbb{Z}_2) \quad (4.24)$$

where $i_1 : C(\Omega) \rightarrow C(\Omega) \rtimes_\sigma \mathbb{Z}_2$ and $i_2 : C(\Omega) \rightarrow C(\Omega) \rtimes_{\varphi\sigma} \mathbb{Z}_2$ are the canonical maps. Note that there is an isomorphism

$$C(\Omega) \rtimes_\sigma \mathbb{Z}_2 \simeq \{f \in C(\Omega, M_2) : f \circ \sigma = \text{Ad } u(f)\} \quad (4.25)$$

induced by the covariant representation (π, u) where $u = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $\pi : C(\Omega) \rightarrow C(\Omega, M_2)$ is given by

$$\pi(f) = \begin{pmatrix} f & 0 \\ 0 & f \circ \sigma \end{pmatrix}.$$

For simplicity of notation we identify $C(\Omega) \rtimes_\sigma \mathbb{Z}_2$ and $\{f \in C(\Omega, M_2) : f \circ \sigma = \text{Ad } u(f)\}$ in the following. There is a map $Tr : C(\Omega) \rtimes_\sigma \mathbb{Z}_2 \rightarrow C(\Omega)$ defined such that

$$Tr(f)(x) = f_{11}(x) + f_{22}(x)$$

when $f \in C(\Omega) \rtimes_{\sigma} \mathbb{Z}_2$. This gives rise to a group homomorphism

$$Tr_* : K_0(C(\Omega) \rtimes_{\sigma} \mathbb{Z}_2) \rightarrow \{g \in C(\Omega, \mathbb{Z}) : g \circ \sigma = g\}.$$

Lemma 4.38. *There is a compact and open set $K \subseteq \Omega$ such that $K \cap \sigma(K) = \emptyset$ and $K \cup \sigma(K) = \Omega$.*

Proof. Let $x \in \Omega$. Since $\sigma(x) \neq x$ there is a clopen subset $U \subseteq \Omega$ such that $U \cap \sigma(U) = \emptyset$. By compactness of Ω we get a finite collection $U_1, U_2, \dots, U_N$ of clopen sets in Ω such that $U_i \cap \sigma(U_i) = \emptyset$ for all i and $\bigcup_{i=1}^N U_i = \Omega$. Set $B_1 = U_1$ and define $B_i, i \geq 2$, recursively, such that

$$B_i = U_i \setminus \left[\bigcup_{j=1}^{i-1} B_j \cup \sigma(B_j) \right].$$

Set $K = B_1 \cup B_2 \cup \dots \cup B_N$. □

Lemma 4.39. *Let K be the clopen set of Lemma 4.38, and define $\beta : C(\Omega) \rtimes_{\sigma} \mathbb{Z}_2 \rightarrow C(K, M_2)$ such that $\beta(f) = f|_K$. Then β is a $*$ -isomorphism.*

Proof. β is clearly an injective $*$ -homomorphism. To see that β is surjective, let $g \in C(K, M_2)$. Define $h : \Omega \rightarrow M_2$ such that

$$h(x) = \begin{cases} g(x), & x \in K \\ ug(\sigma(x))u, & x \in \sigma(K). \end{cases}$$

Then $h \in C(\Omega) \rtimes_{\sigma} \mathbb{Z}_2$ and $\beta(h) = g$. □

Corollary 4.40. *Assume φ is minimal. It follows that*

$$K_1(C(\Omega) \rtimes_{\varphi} \mathbb{Z} \rtimes_{\sigma} \mathbb{Z}_2) = 0.$$

Proof. It follows from Lemma 4.39 that $K_1(C(\Omega) \rtimes_{\sigma} \mathbb{Z}_2) = 0$. For the same reason we have also that $K_1(C(\Omega) \rtimes_{\varphi\sigma} \mathbb{Z}_2) = 0$. The conclusion follows now from the six terms exact sequence of [N]. □

Note that the diagram

$$\begin{array}{ccc} K_0(C(\Omega) \rtimes_{\sigma} \mathbb{Z}_2) & \xrightarrow{Tr_*} & \{f \in C(\Omega, \mathbb{Z}) : f \circ \sigma = f\} \\ \beta_* \downarrow & & \downarrow f \mapsto f|_K \\ K_0(C(K, M_2)) & \xrightarrow{Tr_*} & C(K, \mathbb{Z}) \end{array} \quad (4.26)$$

commutes. It is wellknown and easy to show that the lower Tr_* -map is an isomorphism. It follows in this way from Lemma 4.39 that

$$Tr_* : K_0(C(\Omega) \rtimes_{\sigma} \mathbb{Z}_2) \rightarrow \{f \in C(\Omega, \mathbb{Z}) : f \circ \sigma = f\}$$

is an isomorphism. Similarly,

$$Tr_* : K_0(C(\Omega) \rtimes_{\varphi\sigma} \mathbb{Z}_2) \rightarrow \{f \in C(\Omega, \mathbb{Z}) : f \circ \varphi\sigma = f\}$$

is an isomorphism.

Set $G_\sigma = \{f \in C(\Omega, \mathbb{Z}) : f \circ \sigma = f\}$ and $G_{\varphi\sigma} = \{f \in C(\Omega, \mathbb{Z}) : f \circ \varphi\sigma = f\}$. It follows that $K_0(C(\Omega) \rtimes_\varphi \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2)$ is isomorphic to the cokernel of the map

$$C(\Omega, \mathbb{Z}) \ni h \mapsto (h + h \circ \sigma, h + h \circ \varphi\sigma) \in G_\sigma \oplus G_{\varphi\sigma}.$$

Consider the embedding $\mathbb{Z} \ni z \mapsto (z, z) \in G_\sigma \oplus G_{\varphi\sigma}$ obtained by considering z as a constant function on Ω . If $h \in C(\Omega, \mathbb{Z})$ is such that

$$(i_{1*}(h), i_{2*}(h)) = (h + h \circ \sigma, h + h \circ \varphi\sigma) = (z, z) \quad (4.27)$$

for some $z \in \mathbb{Z}$ we have in particular that $h \circ \sigma = h \circ \varphi\sigma = h \circ \sigma\varphi^{-1}$ which implies that $h \circ \sigma$ and hence also h is constant because φ is minimal. But then (4.27) implies that z is even. This shows that there is an injection

$$t : \mathbb{Z}_2 \rightarrow K_0(C(\Omega) \rtimes_\varphi \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2).$$

By composing the map $G_\sigma \oplus G_{\varphi\sigma} \ni (f, g) \mapsto f - g \in C(\Omega, \mathbb{Z})$ with the quotient map $C(\Omega, \mathbb{Z}) \rightarrow C(\Omega, \mathbb{Z})/(1 - \varphi_*)(C(\Omega, \mathbb{Z}))$ we obtain a map which clearly annihilates elements of the form $(h + h \circ \sigma, h + h \circ \varphi\sigma)$ and hence give rise to a map

$$s : K_0(C(\Omega) \rtimes_\varphi \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2) \rightarrow C(\Omega, \mathbb{Z})/(1 - \varphi_*)(C(\Omega, \mathbb{Z})).$$

Lemma 4.41. *Assume that φ is minimal.*

$$0 \rightarrow \mathbb{Z}_2 \xrightarrow{t} K_0(C(\Omega) \rtimes_\varphi \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2) \xrightarrow{s} (1 + \sigma_*)(C(\Omega, \mathbb{Z})) / (1 - \varphi_*)(C(\Omega, \mathbb{Z})) \rightarrow 0$$

is exact.

Proof. It is obvious that $s \circ t = 0$ so consider an element $(f, g) \in G_\sigma \oplus G_{\varphi\sigma}$ such that $f - g$ represent 0 in $C(\Omega, \mathbb{Z})/(1 - \varphi_*)(C(\Omega, \mathbb{Z}))$. There is then a function $h \in C(\Omega, \mathbb{Z})$ such that

$$f - g = h - \varphi(h). \quad (4.28)$$

Since $f \in G_\sigma, g \in G_{\varphi\sigma}$ we find that

$$f - \sigma(g) = \sigma(h) - \sigma\varphi(h) \quad (4.29)$$

and

$$\varphi(f) - g = \varphi\sigma(h) - \sigma(h). \quad (4.30)$$

Subtracting (4.30) from (4.28) yields that $f - \varphi(f) = h - \varphi(h) - \varphi\sigma(h) + \sigma(h)$ or

$$f - \sigma(h) - h - \varphi(f - \sigma(h) - h) = 0. \quad (4.31)$$

In the same way (4.29) and (4.28) imply that $-g + \sigma(g) = h - \varphi(h) - \sigma(h) + \sigma\varphi(h)$ or (since $\sigma(g) = \varphi^{-1}(g)$),

$$-g + \sigma(h) + \varphi(h) - \varphi^{-1}(-g + \sigma(h) + \varphi(h)) = 0. \quad (4.32)$$

Since φ is minimal we conclude from (4.31) and (4.32) that $f - \sigma(h) - h$ and $-g + \sigma(h) + \varphi(h)$ are constant. Inserting this conclusion into (4.28) we see that the two constants must add to zero, i.e. there is an integer $z \in \mathbb{Z}$ such that i.e. $f - \sigma(h) - h = z$ and $-g + \sigma(h) + \varphi(h) = -z$. Set $k = \sigma(h)$ and note that $(f, g) = (f - k - \sigma(k), g - k - \varphi\sigma(k)) = (z, z)$ in $K_0(C(\Omega) \rtimes_\varphi \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2)$. It follows that the element of $K_0(C(\Omega) \rtimes_\varphi \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2)$ which (f, g) represents is in $t(\mathbb{Z}_2)$.

It remains now only to show that

$$s(K_0(C(\Omega) \rtimes_\varphi \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2)) = (1 + \sigma_*)(C(\Omega, \mathbb{Z})) / (1 - \varphi_*)(C(\Omega, \mathbb{Z})). \quad (4.33)$$

Let $(f, g) \in G_\sigma \oplus G_{\varphi\sigma}$. Let $K \subseteq \Omega$ be the compact open set of Lemma 4.38 and set

$$h_1(x) = \begin{cases} f(x), & x \in K \\ 0, & x \in \sigma(K). \end{cases}$$

Then $h_1 \in C(\Omega, \mathbb{Z})$ and $h_1 + \sigma(h_1) = f$. Similarly there is an $h_2 \in C(\Omega, \mathbb{Z})$ such that $h_2 + \varphi\sigma(h_2) = g$. Since $f - g = (1 + \sigma_*)(h_1 - h_2) + \sigma(h_2) - \varphi\sigma(h_2)$, we see that $f - g \in (1 + \sigma_*)(C(\Omega, \mathbb{Z}))$ modulo $(1 - \varphi_*)(C(\Omega, \mathbb{Z}))$. This proves the inclusion $\subseteq$ in (4.33). Since the reversed inclusion is trivial, the proof is complete. $\square$

Theorem 4.42. *Assume that φ is minimal. Then*

$$K_1(C(\Omega) \rtimes_\alpha D_\infty) = 0$$

and

$$K_0(C(\Omega) \rtimes_\alpha D_\infty) \simeq \mathbb{Z}_2 \oplus (1 + \sigma_*)(C(\Omega, \mathbb{Z})) / (1 - \varphi_*)(C(\Omega, \mathbb{Z})).$$

Proof. In view of Corollary 4.40 and Lemma 4.41 it remains only to show that the $\mathbb{Z}_2$ -subgroup from the extension of Lemma 4.41 is a direct summand. But this follows from a theorem of Kulikov, cf. Theorem 24.5 of [F], since the quotient of the extension, $(1 + \sigma_*)(C(\Omega, \mathbb{Z})) / (1 - \varphi_*)(C(\Omega, \mathbb{Z}))$, is torsion free. $\square$

Remark 4.43. The torsion subgroup $\mathbb{Z}_2$ was overlooked in [BEK]. Its appearance shows that $C(\Omega) \rtimes_\alpha D_\infty$ can fail to be AF when neither σ nor $\varphi\sigma$ has fixed points, a question left open in [BEK]. With this remark we try to give a better description of this torsion element and explain why it disappears when σ or $\varphi\sigma$ has fixed points.

Observe first that the unitary u occurring in (4.25) is unitarily equivalent to $v = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Specifically $wuw^* = v$ where

$$w = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

There is therefore a $*$ -isomorphism $C(\Omega) \rtimes_\sigma \mathbb{Z}_2 \simeq A$ where

$$A = \{f \in C(\Omega, M_2) : f \circ \sigma = \text{Ad } v(f)\}. \quad (4.34)$$

Similarly, there is a $*$ -isomorphism $C(\Omega) \rtimes_{\varphi\sigma} \mathbb{Z}_2 \simeq B$ where

$$B = \{f \in C(\Omega, M_2) : f \circ \varphi\sigma = \text{Ad } v(f)\}. \quad (4.35)$$

Under this identification i_1 and i_2 are given by

$$i_1(h) = \frac{1}{2} \begin{pmatrix} h + h \circ \sigma & h - h \circ \sigma \\ h - h \circ \sigma & h + h \circ \sigma \end{pmatrix},$$

and

$$i_2(h) = \frac{1}{2} \begin{pmatrix} h + h \circ \varphi\sigma & h - h \circ \varphi\sigma \\ h - h \circ \varphi\sigma & h + h \circ \varphi\sigma \end{pmatrix},$$

respectively. The projection

$$p = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

is then an element of both A and B in the obvious way. The element $([p], [p]) \in K_0(A) \oplus K_0(B)$ is not in the range of (i_{1*}, i_{2*}) , for if it was there would have to be a function $h \in C(\Omega, \mathbb{Z})$ such that

$$\text{Tr}_*([p]) = 1 = h + h \circ \sigma = h + h \circ \varphi\sigma. \quad (4.36)$$

The last identity of (4.36) and minimality of φ would force h to be constant which makes the second identity in (4.36) impossible. This shows that $([p], [p])$ represents a non-zero element of $K_0(C(\Omega) \rtimes_{\alpha} D_{\infty})$ when φ is minimal, regardless of whether or not the action α is free. But when it is free the element has order two, because in this case we have the element

$$a = \begin{pmatrix} 0 & 1_K - 1_{\sigma(K)} \\ 0 & 0 \end{pmatrix}$$

in A , where K is the clopen subset of Ω from Lemma 4.38). Since $aa^* = 1_A - p$ and $a^*a = p$ this shows that $[1_A] = 2[p]$ in $K_0(A)$. It follows in the same way that $[1_B] = 2[p]$ in $K_0(B)$. Since $([1_A], [1_B]) = (i_{1*}[1_{C(\Omega)}], i_{2*}[1_{C(\Omega)}])$ we see that $([p], [p])$ represents an element of order 2 in the cokernel of (i_{1*}, i_{2*}) . In contrast, when σ or $\varphi\sigma$ has a fix point there is no substitute for the element a in A or B and the element $([p], [p])$ has in fact infinite order in the cokernel of (i_{1*}, i_{2*}) in this case, as shown in [BEK].

Theorem 4.44. *Let α be a free minimal action of the infinite dihedral group D_{∞} on the Cantor set Ω , generated by the homeomorphisms φ and σ satisfying (4.20) and (4.21). Then*

$$C(\Omega) \rtimes_{\alpha} D_{\infty}$$

is a simple unital AH-algebra with real rank zero and no dimension growth. It is an AT-algebra if and only if φ is not minimal.

Proof. That $C(\Omega) \rtimes_{\alpha} D_{\infty}$ is AT when φ is not minimal follows from Theorem 4.30. When φ is minimal we conclude from Corollary 4.33 and Corollary 4.37 that A has tracial rank zero and satisfies the UCT. It follows then from [Lin4] that A is AH with no dimension growth. That the real rank is zero follows from [Lin1] and [Lin2] because the tracial rank is zero. Finally, it follows from Theorem 4.42 that the algebra is not AT when φ is minimal. $\square$

This ends our excursion into crossed products of the form $C(\Omega) \rtimes_{\alpha} D_{\infty}$ because we have obtained all the information we need for the study of the heteroclinic and homoclinic C^* -algebras of 1-solenoids.

4.6. The inductive limit decomposition of $B_{\bar{h}^{-1}}(\bar{\Gamma})$. We can now put the pieces together and obtain the following

Theorem 4.45. *Let (Γ, h) be a pre-solenoid and $B_{\bar{h}^{-1}}(\bar{\Gamma})$ the heteroclinic algebra of the inverse $\bar{h}^{-1} : \bar{\Gamma} \rightarrow \bar{\Gamma}$ of the 1-solenoid $(\bar{\Gamma}, \bar{h})$. Then $B_{\bar{h}^{-1}}(\bar{\Gamma}) \simeq A \otimes \mathbb{K}$ where A is a unital simple AH-algebra with real rank zero, no dimension growth and a unique trace state.*

Proof. Combine Theorem 4.15 with Proposition 4.27 and Theorem 4.44. $\square$

The K -theory of $B_{\bar{h}^{-1}}(\bar{\Gamma})$ is described in Theorem 4.42 when φ is minimal and in Theorem 4.30 when φ is not minimal. By Lemma 4.17 and Theorem 2.2 φ is minimal if and only if $(\bar{\Gamma}, \bar{h})$ is orientable. This gives us the following conclusion in the same way we obtained Theorem 3.19:

Theorem 4.46. *Let (Γ, h) be an oriented pre-solenoid. Then the heteroclinic algebra $B_{\bar{h}^{-1}}(\bar{\Gamma})$ of $\bar{h}^{-1} : \bar{\Gamma} \rightarrow \bar{\Gamma}$ is a simple stable AT-algebra of real rank zero.*

Example 4.47. The K_0 -groups of $K_0(B_{\bar{h}^{-1}}(\bar{\Gamma}))$ agrees with the first Čech cohomology group of $\bar{\Gamma}$ when (Γ, h) is oriented. This follows from [Y2] and [Y4]. There is even a natural way to define an order on the first Čech cohomology group which corresponds to the order on $K_0(B_{\bar{h}^{-1}}(\bar{\Gamma}))$.

I don't know if $K_0(B_{\bar{h}}(\bar{\Gamma}))$ has an interpretation in terms of known invariants for 1-solenoids, even in the orientable case. In many cases the two groups, $K_0(B_{\bar{h}^{-1}}(\bar{\Gamma}))$ and $K_0(B_{\bar{h}}(\bar{\Gamma}))$ are isomorphic as dimension groups and it follows from the results in this paper that $B_{\bar{h}^{-1}}(\bar{\Gamma})$ and $B_{\bar{h}}(\bar{\Gamma})$ are then isomorphic. This happens for example for the 1-solenoids of Remark 2.1. In fact, it is not so easy to come up with an example of an oriented 1-solenoid for which $B_{\bar{h}^{-1}}(\bar{\Gamma}) \neq B_{\bar{h}}(\bar{\Gamma})$, but here is one: In the setting of Remark 2.1 let $h : \Gamma \rightarrow \Gamma$ be given by the wrapping rule

$$a \mapsto a^{65}b^7, \quad b \mapsto a^{24}b^{67}.$$

By Corollary 3.7 of [Y4] $K_0(B_{\bar{h}^{-1}}(\bar{\Gamma}))$ is the stationary dimension group given by the matrix

$$A = \begin{pmatrix} 65 & 7 \\ 24 & 67 \end{pmatrix}.$$

This group is not isomorphic to $K_0(B_{\bar{h}}(\bar{\Gamma}))$, even when we ignore the order. Indeed, by using the description of $B_{\bar{h}}(\bar{\Gamma})$ given in Section 3 it can be shown that $K_0(B_{\bar{h}}(\bar{\Gamma}))$ is the stationary dimension group corresponding to the *transpose* $A^t = \begin{pmatrix} 65 & 24 \\ 7 & 67 \end{pmatrix}$ of A . These dimension groups are not even isomorphic as groups, cf. Example 3.6 of [BJKR] and Appendix C of [Th1].

5. THE HOMOCLINIC ALGEBRA OF A 1-SOLENOID

By using, for the first time, that 1-solenoids are Smale spaces we can combine the preceding results with [Pu] to obtain the following

Theorem 5.1. *Let (Γ, h) be a pre-solenoid. The homoclinic C^* -algebra $A_{\bar{h}}(\bar{\Gamma})$ of the 1-solenoid $(\bar{\Gamma}, \bar{h})$ is a simple unital AH-algebra of real rank zero with no dimension growth and a unique trace state.*

Proof. By Theorem 3.1 of [Pu] $A_{\bar{h}}(\bar{\Gamma})$ is stably isomorphic to $B_{\bar{h}}(\bar{\Gamma}) \otimes B_{\bar{h}^{-1}}(\bar{\Gamma})$. It follows then from Theorem 4.45 and Theorem 3.14 that $A_{\bar{h}}(\bar{\Gamma})$ is stably isomorphic to a tensor-product $D_1 \otimes D_2$ where D_1 and D_2 are simple unital AH-algebras of real rank zero with no dimension growth and a unique trace state. It is standard to deduce from this that $D_1 \otimes D_2$ itself is a simple unital AH-algebra with no dimension growth and a unique trace state. To see that $D_1 \otimes D_2$ has real rank zero note that AH-algebras without dimension growth are inductive limits of recursive sub-homogeneous algebras in the sense of [Ph3]. It follows then from Proposition 1.10 and Theorem 4.2 of [Ph3] that $D_1 \otimes D_2$ has real rank zero. $\square$

Since $A_{\bar{h}}(\bar{\Gamma})$ is stably isomorphic to the tensor product $B_{\bar{h}^{-1}}(\bar{\Gamma}) \otimes B_{\bar{h}}(\bar{\Gamma})$ we can find the K-groups of $A_{\bar{h}}(\bar{\Gamma})$ by using the results from Section 3 and Section 4 in combination with the Künneth theorem, [Bl].

By combining Theorem 4.46 and Theorem 3.19 we obtain the following

Theorem 5.2. *Let (Γ, h) be an oriented pre-solenoid. Then the homoclinic algebra $A_{\overline{h}}(\overline{\Gamma})$ is a simple unital AT-algebra of real rank zero with a unique trace state.*

REFERENCES

- [AO] J. M. Aarts and L. Oversteegen, *Matchbox manifolds*, Continua (Cincinnati, OH, 1994), 3-14, Lecture Notes in Pure and Appl. Math., 170, Dekker, New York, 1995.
- [BEK] O. Bratteli, D.E. Evans and A. Kishimoto, *Crossed products of totally disconnected spaces by $\mathbb{Z}_2 * \mathbb{Z}_2$* , Ergod. Th. & Dynam. Sys. **13** (1993), 445-484.
- [Bl] B. Blackadar, *K-theory for Operator Algebras*, Springer Verlag, New York, 1986.
- [BJKR] O. Bratteli, P. E. T. Jorgensen, K.H. Kim and F. Roush *Computation of isomorphism invariants for stationary dimension groups*, Ergod. Th. & Dynam. Sys. **22** (2002), 99-127.
- [Br] L. Brown, *Stable isomorphism of hereditary subalgebras of C^* -algebras*, Pac. J. Math. **71** (1977), 335-348.
- [BP] L. Brown and G.K. Pedersen, *C^* -algebras of real rank zero*, J. Funct. Anal. **99** (1991), 132-149.
- [CuK] J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Invent. Math. **56** (1980), 25-268.
- [Ef] E. Effros, *Dimensions and C^* -algebras*, CBMS Regional Conf. Ser. in Math., no. 46, Amer. Math. Soc., Providence, 1981.
- [Ell1] G. Elliott, *A classification of certain simple C^* -algebras, II*, J. Ramanujan Math. Soc. **12** (1997), 97-134.
- [Ell2] ———, *On the classification of C^* -algebras of real rank zero*, J. Reine Angew. Math. **443** (1993), 179-219.
- [EG] G. Elliott and G. Gong, *On the classification of C^* -algebras of real rank zero, II*, Ann. Math. **144** (1996), 497-610.
- [EGL] G. Elliott, G. Gong and L. Li, *On the classification of simple inductive limit C^* -algebras. II. The isomorphism theorem*, Invent. Math. **168** (2007), no. 2, 249-320.
- [F] R. Fokink, *The structure of trajectories*, Ph.D. Thesis, Technische Universiteit te Delft, 1991.
- [F] L. Fuchs, *Abelian groups*, Pergamon Press, Oxford, London, New York, Paris, 1960.
- [GPS1] T. Giordano, I. Putnam and C. Skau, *Topological orbit equivalence and C^* -crossed products*, Journal für die reine und angewandte Mathematik **469** (1995), 51-111.
- [GPS2] ———, *Affable equivalence relations and orbit structure of Cantor dynamical systems*, Ergodic Theory Dynam. Systems **24** (2004), no. 2, 441-475.
- [HPS] R.H. Herman, I.F. Putnam and C.F. Skau, *Ordered Bratteli diagrams, dimension groups and topological dynamics*, Internat. J. Math. **3** (1992), 827-864.
- [HR] J. v. B. Hjelmberg, M. Rørdam, *On stability of C^* -algebras*, J. Funct. Anal. **155** (1998), 152-170.
- [Kr1] W. Krieger, *On a dimension for a class of homeomorphism groups*, Math. Ann. **252** (1979/80), 87-95.
- [Kr2] ———, *On dimension functions and topological Markov chains*, Invent. Math. **56** (1980), 239-250.
- [Lin1] H. Lin, *Tracially AF C^* -algebras*, Trans. Amer. Math. Soc. **353** (2001), 693-722.
- [Lin2] ———, *The tracial topological rank of C^* -algebras*, Proc. London Math. Soc. **83** (2001), 199-234.
- [Lin3] ———, *Traces and simple C^* -algebras with tracial topological rank zero*, J. Reine angew. Math. **568** (2004), 99-137.
- [Lin4] ———, *Classification of simple C^* -algebras of tracial rank zero*, Duke Math. J. **125** (2004), 91-119.
- [MRW] P. Muhly, J. Renault and D. Williams, *Equivalence and isomorphism for groupoid C^* -algebras*, J. Oper. Th. **17** (1987), 3-22.
- [N] T. Natsume, *On $K_*(C^*(SL_2(\mathbb{Z})))$* , J. Operator Theory **13** (1985), 103-118.
- [OP] H. Osaka and N. C. Phillips, *Crossed products by finite groups with the Rokhlin property*, Preprint (2007), arXiv:math/0704.3651v [math.OA]

- [Ph1] N. C. Phillips, *Crossed Products of the Cantor Set by Free Minimal Actions of $\mathbb{Z}^d$* , Comm. Math. Phys. **256** (2005), 1–42.
- [Ph2] ———, *Every simple higher dimensional noncommutative torus is an AT algebra*, Preprint, September, 2006.
- [Ph3] ———, *Real rank and property (SP) for direct limits of recursive subhomogeneous algebras*, Trans. Amer. Math. Soc., to appear.
- [Ph4] ———, *The tracial Rokhlin property for actions of finite groups on C^* -algebras*, Preprint (2006), arXiv:math/0609782v1 [math.OA]
- [Ph5] ———, *Finite cyclic group actions with the tracial Rokhlin property*, Preprint (2006), arXiv:math/0609785v1 [math.OA]
- [Pu] I. Putnam, *C^* -algebras from Smale spaces*, Can. J. Math. **48** (1996), 175–195.
- [Re] J. Renault, *A Groupoid Approach to C^* -algebras*, LNM 793, Springer Verlag, Berlin, Heidelberg, New York, 1980.
- [Ru1] D. Ruelle, *Thermodynamic Formalism*, Encyclopedia of Mathematics and its Applications 5, Addison-Wesley, Reading, Ma. 1978.
- [Ru2] ———, *Non-commutative algebras for hyperbolic diffeomorphisms*, Invent. Math. **93** (1988), 1–13.
- [Th1] K. Thomsen, *C^* -algebras of homoclinic and heteroclinic structure in expansive dynamics*, Preprint, IMF Aarhus University, 2007.
- [Th2] ———, *Traces, Unitary Characters and Crossed Products by $\mathbb{Z}$* , Publ. RIMS **31**(1995), 1011–1029.
- [Th3] ———, *On the KK-theory and the E-theory of amalgamated free products of C^* -algebras*, J. Func. Anal. **201** (2003), 30–56.
- [Y1] I. Yi, *Canonical symbolic dynamics for one-dimensional generalized solenoids*, Trans. Amer. Math. Soc. **353** (2001), 3741–3767.
- [Y2] ———, *Ordered group invariants for one-dimensional spaces*, Fund. Math. **170** (2001), 267–286.
- [Y3] ———, *Ordered group invariants for nonorientable one-dimensional generalized solenoids*, Proc. Amer. Math. Soc. **131** (2003), 1273–1282.
- [Y4] ———, *K-theory of C^* -algebras from one-dimensional generalized solenoids*, J. Operator Theory **50** (2003), 283–295.
- [Y5] ———, *Bratteli-Vershik systems for one-dimensional generalized solenoids*, Houston J. Math. **30** (2004), 691–704.
- [Wi1] R.F. Williams, *One-dimensional non-wandering sets*, Topology **6** (1967), 473–487.
- [Wi2] ———, *Classification of one dimensional attractors*, Proceedings of Symposia in Pure Mathematics **14** (1970), 341–361.

E-mail address: matkt@imf.au.dk

INSTITUT FOR MATEMATISKE FAG, NY MUNKEGADE, 8000 AARHUS C, DENMARK